

REAL K3 SURFACES WITH NON-SYMPLECTIC INVOLUTION AND APPLICATIONS. II

VIACHESLAV V. NIKULIN AND SACHIKO SAITO

ABSTRACT. We consider real forms of relatively minimal rational surfaces $\mathbb{F}_m$. Connected components of moduli of real non-singular curves in $|-2K_{\mathbb{F}_m}|$ had been classified recently for $m = 0, 1, 4$ in [19]. Applying similar methods, here we fill the gap for $m = 2$ and $m = 3$ to complete similar classification for any $0 \leq m \leq 4$, when $|-2K_{\mathbb{F}_m}|$ is reduced.

The case of $\mathbb{F}_2$ is especially remarkable and classical (quadratic cone in $\mathbb{P}^3$). As an application, we finished classification of connected components of moduli of real hyper-elliptically polarized K3 surfaces and their deformations to real polarized K3 surfaces started in [19] and [18]. This could be important in some questions because real hyper-elliptically polarized K3 surfaces can be constructed explicitly.

CONTENTS

1. Introduction	2
2. Connected components of moduli of real non-singular curves in $ -2K_{\mathbb{F}_3} $	3
2.1. Reduction to real K3 surfaces with non-symplectic involution of type $(3, 1, 1)$	4
2.2. Enumeration of connected components of moduli of real K3 surfaces with non-symplectic involution	
2.3. Geometric interpretation of the invariants (2.4) in terms of K3	9
2.4. Geometric interpretation of the invariants (2.4) in terms of $A \in -2K_{\mathbb{F}_3} $	10
3. Connected components of moduli of real non-singular curves in $ -2K_{\mathbb{F}_2} $	13
3.1. Reduction to real K3 surfaces with degenerate non-symplectic involution of type $(2, 2, 0)$	14
3.2. Enumeration of connected components of moduli of real K3 surfaces with degenerate non-symplectic involution	
3.3. Geometric interpretation of the invariants (3.3) in terms of K3	20
3.4. Geometric interpretation of the invariants (3.3) in terms of $A \in -2K_{\mathbb{F}_2} $	21
3.5. Application: Connected components of moduli of real non-singular degree 8 curves on a quadratic cone	
3.6. Application: Hyperboloid deformations of $A \in -2K_{\mathbb{F}_2} $ to $A \in -2K_{\mathbb{F}_0} $.	25
3.7. Application: Ellipsoid deformations of $A \in -2K_{\mathbb{F}_2} $ to $A \in -2K_{\mathbb{F}_0} $.	26
3.8. Application: Classification of connected components of moduli of real hyper-elliptically polarized K3 surfaces	
3.9. Application: Real polarized K3 surfaces as deformations of real hyper-elliptically polarized K3 surfaces	
References	32

2000 *Mathematics Subject Classification.* 14H45, 14J26, 14J28, 14P25.

Key words and phrases. real K3 surface, involution, real Enriques surface, real rational surface, real algebraic curve, ellipsoid, hyperboloid.

1. INTRODUCTION

In [19] classification of connected components of moduli of real K3 surfaces with non-symplectic involution had been considered. As an application, classification of connected components of moduli of real non-singular curves $A \in |-2K_{\mathbb{F}_m}|$ had been obtained for $m = 0, 1, 4$ where $\mathbb{F}_m$ is a relatively minimal rational surface having exceptional section with square $-m$. Here a K3 surface with non-symplectic involution appears as the double coverings of $\mathbb{F}_m$ ramified in A with the involution of the double covering.

The main purpose of this paper is to fill the gap, and to get similar classification for $m = 2$ and $m = 3$.

Case of $\mathbb{F}_3$ (see Sect. 2) is very similar to cases $m = 0, 1, 4$ considered in [19] because in this case we obtain K3 surfaces with non-degenerate non-symplectic involution. This non-degenerate case had been treated in [19] in general. Remarkably, enumeration of the connected components for $\mathbb{F}_3$ case is equivalent to classification of some special integral polarized involutions classified in [15] in general. Thus, direct application of results of [15] and [19] solves the problem in this case.

The case of $\mathbb{F}_3$ is related with some very classical problems.

Firstly, K3 surfaces of this case can be also considered as K3 surfaces with fixed elliptic pencil and its section such that the elliptic pencil has no reducible fibres except one having the type corresponding to the extended Dynkin diagram $\tilde{A}_1$ or $\tilde{A}_2$. The inverse map of the group law of the elliptic pencil with the section gives then the non-symplectic involution. Thus, we classify connected components of moduli of such real K3 surfaces. Similarly, the case of $\mathbb{F}_4$ which had been treated in [19] is equivalent to K3 surfaces having fixed elliptic pencil with section and no reducible fibres.

Secondly, the case of non-singular curves $A \in |-2K_{\mathbb{F}_3}|$ is equivalent to curves $A \in |-2K_{\mathbb{F}_4}|$ having no singular points except one quadratic (possibly degenerate) singular point. Thus, we describe connected components of their moduli over $\mathbb{R}$ too. See Remark 8.

Thirdly, non-singular curves $A \in |-2K_{\mathbb{F}_3}|$ are equivalent to non-singular curves $A_1 \in |-2K_Z|$ where Z is a del Pezzo surface having no singular points except one log-terminal singular point of index two having the type K_1 , moreover, the genus $g(A_1) = 9$ (see classification of log del Pezzo surfaces of index ≤ 2 in [1]). Thus, we classify connected components of moduli of real non-singular curves $A_1 \in |-2K_Z|$ of such real del Pezzo surfaces Z . See Remark 9 for details.

The case of $\mathbb{F}_2$ (see Sect. 3) when we consider connected components of moduli of real non-singular curves $A \in |-2K_{\mathbb{F}_2}|$ is especially remarkable. It gives K3 surfaces with degenerate non-symplectic involution since $\mathbb{F}_2$ has exceptional curves with square (-2) . Thus, we cannot directly apply results of [19] where only non-degenerate case had been treated. Fortunately, all main results of [19] survive in this case, and we show that the connected component of moduli is also determined by the action (L, τ, φ) of τ (holomorphic non-symplectic involution) and φ (anti-holomorphic involution) on the homology lattice $L = H_2(X(\mathbb{C}); \mathbb{Z})$ of a K3 surface X from this connected component. It is clear that only in very special cases of degenerate non-symplectic involutions on K3 similar result is valid. So, we are very lucky. It would be interesting to describe all similar nice cases in general, see Remark 11. See [1]

about classification of K3 surfaces over $\mathbb{C}$ with degenerate (i.e. arbitrary) non-symplectic involution.

Remarkably, enumeration of the connected components of moduli in the case of $\mathbb{F}_2$ is also equivalent to classification of some special integral polarized involutions. Thus, again application of results of [15] and [19] solves the problem.

The case of $\mathbb{F}_2$ is related with several classical problems.

Firstly, $\mathbb{F}_2$ with contracted exceptional section is a quadratic cone in $\mathbb{P}^3$. Thus, we solve very classical problem of enumeration of connected components of moduli of non-singular degree 8 curves on the quadratic cone in 3-dimensional space (i. e. non-singular sections of the cone by degree four forms). See Sect. 3.5.

Deforming the cone to hyperboloid or ellipsoid, we obtain bidegree (4,4) non-singular curves on hyperboloid or ellipsoid (the case of $\mathbb{F}_0$). Combining our results here and results of [19] we completely classify these deformations. This could be important in some very practical problems. See Sects 3.6 and 3.7.

Let us consider hyper-elliptically polarized K3 surfaces (X, P) such that the linear system $|P|$ is hyper-elliptic and gives then a double covering $|P| : X \rightarrow Y$. The case of $Y \cong \mathbb{F}_2$ gives *hyper-elliptically polarized K3 surfaces (X, P) of type $\mathbb{F}_2$* . In this case, $|P|$ is double covering of $\mathbb{F}_2$ ramified in a non-singular curve $A \in |-2K_{\mathbb{F}_2}|$. Thus, applying our results, we obtain classification of connected components of moduli of real hyper-elliptically polarized K3 surfaces of type $\mathbb{F}_2$. This finishes classification of connected components of moduli of real hyper-elliptically polarized K3 surfaces since for all other types similar classification had been obtained in [19] (see also [18]). See Sect. 3.8.

In Sect. 3.9, applying our results about $\mathbb{F}_2$ and results of [19] and [18], we classify deformations of real hyper-elliptically polarized K3 surfaces of type $\mathbb{F}_2$ to real polarized K3 surfaces. We completely describe all real polarized K3 surfaces which can be obtained by such deformation. Moreover, we enumerate all types of these deformations. Similar classification for all other types of real hyper-elliptically polarized K3 surfaces had been obtained in [19] and [18]. Thus, here we finalized these results.

All our results here and in [19], [18] are based on Global Torelli Theorem [20] and surjectivity of Torelli map [14] for K3 surfaces, and some geometric and arithmetic considerations. Thus, our results are mainly existence and uniqueness results. It would be very interesting to construct exactly all representatives of connected components of moduli which we considered and classified. Our results about deformations reduce everything to construction of representatives of connected components of moduli of real non-singular curves $A \in |-2K_{\mathbb{F}_m}|$, $0 \leq m \leq 4$, or degree 6 curves in $\mathbb{P}^2$. In many cases they are known by direct construction. E. g. see [7], [6], [8], [21], [24], [26], [27], [28]. It would be very interesting to have such constructions for all connected components of moduli. It will make our results effective and useful for some questions which don't depend on deformation. This construction is surely possible because all our connected components of moduli are exactly labelled by some natural invariants, and number of the connected components is bounded.

2. CONNECTED COMPONENTS OF MODULI OF REAL NON-SINGULAR CURVES IN $|-2K_{\mathbb{F}_3}|$

We always consider algebraic varieties over $\mathbb{C}$. If we say “real”, we consider algebraic varieties which are defined over $\mathbb{R}$.

We denote by $\mathbb{F}_m$, $m \geq 0$, a relatively minimal rational surface with exceptional (if $m \geq 1$) section $s \cong \mathbb{P}^1$ having $s^2 = -m$. If $m = 0$, then $\mathbb{F}_0 = \mathbb{P}^1 \times \mathbb{P}^1$. If $m > 0$, then $\mathbb{F}_m$ has a unique rational pencil with the section s . We denote by c a fibre of the rational pencil. The canonical class $K_{\mathbb{F}_m} = (-m - 2)c - 2s$ and $-2K_{\mathbb{F}_m} = (2m + 4)c + 4s$.

If $0 \leq m \leq 2$, then a general element of the complete linear system $| - 2K_{\mathbb{F}_m} |$ is a non-singular curve.

If $3 \leq m \leq 4$, then $| - 2K_{\mathbb{F}_m} | = |(2m + 4)c + 3s| + s$ has the fixed component s where the general element $A_1 \in |(2m + 4)c + 3s|$ is non-singular. If $m = 3$, then $A_1 \cdot s = 1$, and $A = A_1 + s \in | - 2K_{\mathbb{F}_3} |$ is union of two non-singular curves which intersect transversally in one point (for $m = 3$, we then also call A non-singular). If $m = 4$, then $A_1 \cdot s = 0$, and the whole curve $A = A_1 + s \in | - 2K_{\mathbb{F}_4} |$ is non-singular.

If $m > 4$, then $| - 2K_{\mathbb{F}_m} | = |(2m + 4)c + 2s| + 2s$ has a non-reduced fixed component $2s$.

Thus, cases $0 \leq m \leq 4$ are characterized by the property that a general element of the complete linear system $| - 2K_{\mathbb{F}_m} |$ is reduced.

Connected components of moduli of real non-singular curves in $| - 2K_{\mathbb{F}_m} |$ (for real $\mathbb{F}_m$) were classified in [19] for $m = 0, 1, 4$. For $m = 1$ it can be, in principle, deduced from [10], [11] where much more delicate classification had been considered. On the other hand, our classification in [19] gives *natural invariants* of connected components of moduli of real non-singular curves in $| - 2K_{\mathbb{F}_1} |$.

In this paper, we want to complete this classification for reduced $| - 2K_{\mathbb{F}_m} |$ (i. e. for $0 \leq m \leq 4$) considering connected components of moduli of real non-singular curves in $| - 2K_{\mathbb{F}_3} |$ and $| - 2K_{\mathbb{F}_2} |$.

In this section, we consider the case of $| - 2K_{\mathbb{F}_3} |$ which is very similar to $| - 2K_{\mathbb{F}_m} |$ for cases $m = 0, 1, 4$ considered in [19]. The case $| - 2K_{\mathbb{F}_3} |$ is completely covered by the general theory which had been developed in [19]. One needs only to perform necessary calculations and geometric observations corresponding to this case.

The case of $| - 2K_{\mathbb{F}_2} |$ is more delicate. It requires some changes comparing to [19], and it will be considered in the next Sect. 3.

2.1. Reduction to real K3 surfaces with non-symplectic involution of type $(3, 1, 1)$.

Here we consider $\mathbb{F}_3$. Like above, c denotes the fibre of the rational pencil of $\mathbb{F}_3$, and s its exceptional section. We have $c^2 = 0$, $s^2 = -3$ and $c \cdot s = 1$.

Let $A \in | - 2K_{\mathbb{F}_3} |$ be a non-singular curve, i. e. $A = A_1 + s$ where A_1 is a non-singular curve of the genus 9. Then $A_1 \cdot s = 1$, and A_1 intersects s transversally in one point $A_1 \cap s$. Let f be the fibre of the rational pencil of $\mathbb{F}_3$ over this point.

Let $\sigma : Y \rightarrow \mathbb{F}_3$ be the blow-up at the point $A_1 \cap s$ and e the exceptional curve of the blow-up. The Y is a rational surface with Picard number 3. We denote by A_0 the proper pre-image of s , and by the same letters A_1 , f the proper pre-images of A_1 and f on Y . On Y , we have $e^2 = f^2 = -1$, $A_0^2 = -4$ and $A_0 \cdot e = e \cdot f = 1$, and $A_0 \cdot f = 0$. Moreover, $A_1 \cdot A_0 = 0$, $A_1 \cdot e = 1$ and $A_1 \cdot f = 2$.

The proper pre-image $A = A_1 + A_0 \in | - 2K_Y |$ of A is a non-singular curve. Let $\pi : X \rightarrow Y$ be the double covering of Y ramified in $A \subset Y$. The X is a K3 surface with the non-symplectic involution τ of the double covering. Then $A = X^\tau$ is identified with the fixed points set of the involution, and $Y = X/\{1, \tau\}$ is the quotient space. We denote $E = \pi^{-1}(e)$, $F = \pi^{-1}(f)$, and we identify $\pi^{-1}(A_0)$ with A_0 . We have on X that $A_0^2 = E^2 = F^2 = -2$

and $A_0 \cdot E = 1$, $A_0 \cdot F = 0$ and $E \cdot F = 2$. The curves A_0 and E are non-singular rational, i. e. they are isomorphic to $\mathbb{P}^1$. The curve F is also non-singular rational if A_1 intersects f in two distinct points. If A_1 touches f , then $F = F' + F''$ is union of two non-singular rational curves which are conjugate by τ , and $F' \cdot F'' = 1$.

The determinant of the Gram matrix of A_0 , E and F is 2. It follows that A_0 , E and F generate a primitive 3-dimensional sublattice S in the Picard lattice S_X of X which is the eigenvalue 1 sublattice $S = S_X^\tau = H_2(X(\mathbb{C}); \mathbb{Z})^\tau$ of the involution τ (since Y has the Picard number 3). Thus, the lattice $S = S_X^\tau$ is a hyperbolic lattice of the rank $r(S) = 3$. The discriminant group $S^*/S \cong (\mathbb{Z}/2\mathbb{Z})^1$ is a 2-elementary group of the rank $a(S) = 1$. The element $cl(F)/2 \in S^*$ has $(F/2)^2 = -2/4 \notin \mathbb{Z}$. Thus, the 2-elementary lattice S has the invariant $\delta(S) = 1$ (it would be 0 if any element $x^* \in S^*$ has $(x^*)^2 \in \mathbb{Z}$). Thus, finally, the 2-elementary hyperbolic lattice S has the invariants $(r(S), a(S), \delta(S)) = (3, 1, 1)$ which are the main invariants of the K3 surface with non-symplectic involution (X, τ) .

Vice versa, by [1], any K3 surface with non-symplectic involution (X, τ) having the lattice $S = S_X^\tau$ with the same invariants $(r(S), a(S), \delta(S)) = (3, 1, 1)$ can be obtained in the same way as above from a non-singular curve $A_1 + s \in |-2K_{\mathbb{F}_3}|$. Moreover, all exceptional curves (i. e. irreducible with negative self-intersection) on $Y = X/\{1, \tau\}$ are exactly the curves A_0 , e and f . Thus, one obtains the non-singular pair $A_1 + s \in |-2K_{\mathbb{F}_3}|$ from (X, τ) in a unique way by contracting e and taking the image of $A = X^\tau$ which consists of disjoint union of a curve A_1 of genus 9 and $A_0 \cong \mathbb{P}^1$. Thus, pairs (X, τ) with the invariants $(3, 1, 1)$ and non-singular $A_1 + s \in |-2K_{\mathbb{F}_3}|$ are in one to one correspondence. In particular, they have the same moduli space and its connected components.

We mention that K3 surfaces with non-symplectic involution (X, τ) of type $(3, 1, 1)$ give a very important class of K3 surfaces. They are K3 surfaces with elliptic pencil $|E + F|$ (it is pre-image of the rational pencil of $\mathbb{F}_3$), its fixed section A_0 and a reducible fibre $E + F$ satisfying conditions (a), (b) and (c) from (2.1) below:

(2.1)

- (a) E is irreducible non-singular rational and $E \cdot A_0 = 1$;
- (b) F is either irreducible non-singular rational or is sum of two irreducible non-singular rational curves, and $E \cdot F = 2$ (i. e. the degenerate fibre corresponds to the extended root system $\tilde{A}_1$ or $\tilde{A}_2$);
- (c) the elliptic pencil does not have other reducible fibres.

Then τ is the inverse map of the group law of the elliptic pencil with the section. It is a non-symplectic involution of type $(3, 1, 1)$.

Similarly, the case of non-singular curves $A \in |-2K_{\mathbb{F}_4}|$ which had been treated in [19] is equivalent to K3 surfaces with fixed elliptic pencil with section and without reducible fibres (in this case, $(r(S), a(S), \delta(S)) = (2, 0, 0)$).

Now let us assume that $\mathbb{F}_3$ and $A = A_1 + s \in |-2K_{\mathbb{F}_3}|$ are real. Then the point $A_1 \cap s$ is real and $\mathbb{F}_3(\mathbb{R})$ is non-empty. Then $Y(\mathbb{R})$ is non-empty and the corresponding anti-holomorphic involution θ of Y can be lifted to an anti-holomorphic involution φ of the K3 surface X . Thus, we obtain a real K3 surface with non-symplectic involution (X, τ, φ) , i. e. τ and φ commute. The lift φ is unique up to replacement by the *related anti-holomorphic involution* $\tilde{\varphi} = \varphi\tau$.

This choice is defined by the image $\pi(X_\varphi(\mathbb{R})) = A^+(\varphi) \subset Y(\mathbb{R})$ where $X_\varphi(\mathbb{R}) = X(\mathbb{C})^\varphi$ is the set of real points of the real K3 surface (X, φ) . Here $A^+(\varphi)$ is called the *positive curve of A*. If one takes the related real structure $\tilde{\varphi} = \varphi\tau$ on X , then the positive curve becomes $\pi(X_{\tilde{\varphi}}(\mathbb{R})) = A^+(\tilde{\varphi}) = A^-(\varphi) = \overline{Y(\mathbb{R}) - A^+(\varphi)}$. We have $A^+(\varphi) \cap A^+(\tilde{\varphi}) = A(\mathbb{R})$.

Thus, classifying connected components of moduli of real K3 surfaces with non-symplectic involution (X, τ, φ) of type $(3, 1, 1)$, we actually solve more delicate problem of classification of connected components of moduli of positive real non-singular curves A^+ where $A \in |-2K_{\mathbb{F}_3}|$ is non-singular. Here A^+ is defined by a choice of one of two parts of $\mathbb{F}_3(\mathbb{R}) - A(\mathbb{R})$ where $A^+ \cap (\mathbb{F}_3(\mathbb{R}) - A^+) = A(\mathbb{R})$.

2.2. Enumeration of connected components of moduli of real K3 surfaces with non-symplectic involution of type $(3, 1, 1)$. Here we describe connected components of moduli of real K3 surfaces with non-symplectic involution of type $(3, 1, 1)$.

For any (X, τ) of type $(3, 1, 1)$, the surface Y has no exceptional curves with square (-2) . Thus, (X, τ) is always non-degenerate in the sense of ([19], Sect. 2.1). In this case, a connected component of moduli of real K3 surfaces with non-symplectic involution (X, τ, φ) is determined by the isomorphism class of the action (L, τ, φ) of τ and φ on the homology lattice $L = H_2(X(\mathbb{C}); \mathbb{Z})$ with intersection pairing ([19], Theorem 1). Here L is an even unimodular lattice of signature $(3, 19)$.

Since A_0 , e and f are all exceptional curves on Y and they have different geometry on Y , the anti-holomorphic involution θ on Y should send each of these three curves to itself changing its orientation. It follows that $\varphi|_S = \varphi|L^\tau = -1$. Moreover, the eigenvalue 1 part L^φ of φ in L must be hyperbolic (it has exactly one positive square). By ([19], Theorem 1), any isomorphism class of triplets

$$(2.2) \quad (L, \tau, \varphi)$$

with these properties is possible. These isomorphism classes are in one-to-one correspondence with connected components of moduli. Moreover by ([19], Theorem 14 and Proposition 15) the isomorphism classes are determined by their genus invariants.

The genus invariants can be enumerated using general results of [17] which are valid for any S (see also their adaptation to the particular case of K3 with non-symplectic involution in [19]). On the other hand, our case is very simple, and it can be also reduced to invariants of integral polarized involutions (see Sect. 3 in [15]). In this case, all necessary calculations are already done in Theorem 3.4.3 of [15]. We prefer the second opportunity.

The lattice $S = U \oplus \mathbb{Z}cl(F)$ where $U = \mathbb{Z}cl(A_0) + \mathbb{Z}(cl(E) + cl(F))$ is the hyperbolic even unimodular lattice of signature $(1, 1)$, and $\mathbb{Z}cl(F)$ is isomorphic to $\langle -2 \rangle$ since $F^2 = -2$. Here $\oplus$ means orthogonal sum of lattices, and $\langle M \rangle$ a lattice with the symmetric matrix M . Since $\tau|_U = 1$ and $\varphi|_U = -1$, a triplet (L, τ, φ) defines an integral polarized involution

$$(2.3) \quad (L_1, \varphi_1, cl(F)).$$

Here L_1 is the orthogonal complement to U in L (thus, it is an even unimodular lattice of signature $(2, 18)$, $\varphi_1 = \varphi|_{L_1}$ and $L_1^{\varphi_1}$ is hyperbolic, moreover, $\varphi_1(cl(F)) = -cl(F)$ and $F^2 = -2$).

Taking orthogonal sum of (2.3) with U , one obtains the triplet (2.2). Thus, their isomorphism classes are in one to one correspondence.

Genus invariants of integral polarized involutions (2.3) (and then (2.2)) were completely classified in ([15], Theorem 3.4.3). Thus, they enumerate the connected components of moduli. Below we present these results. (One should multiply by (-1) the pairing in L_1 and apply Theorem 3.4.3 in [15] to $l_{(+)} = 18, l_{(-)} = 2, t_{(-)} = 1, t_{(+)} = r - 1$ and $n = 2$.)

The complete genus and isomorphism class invariants of (2.3) (and then (2.2)) are

$$(2.4) \quad (r, a, \delta_\varphi; \delta_F, \delta_{\varphi F}).$$

Here $r = \text{rk } L^\varphi \in \mathbb{N}$; $((L^\varphi)^*/L^\varphi) = (\mathbb{Z}/2\mathbb{Z})^a$ where $a \geq 0$ is an integer; $\delta_\varphi \in \{0, 1\}$ is equal to 0 if and only if $x \cdot \varphi(x) \equiv 0 \pmod{2}$ for any $x \in L$. They are complete invariants of the corresponding pair (L, φ) . Here $\delta_F \in \{0, 1\}$ is equal to 0 if and only if $F \cdot L_\varphi \equiv 0 \pmod{2}$; here $\delta_{\varphi F} \in \{0, 1\}$ is equal to 0 if and only if $x \cdot \varphi(x) \equiv x \cdot F$ for any $x \in L$. Here and in what follows, we always denote by L^φ and L_φ the eigenvalue 1 and -1 parts respectively of the action of an involution φ on a module L .

The invariants (2.4) must satisfy the following relations which are sufficient and necessary for existence of a triplet (2.3) or the corresponding triplet (2.2) with these invariants. Thus, these relations enumerate connected components of moduli.

0. Conditions on (r, a, δ_φ) :

- (1) $1 \leq r \leq 18, 0 \leq a \leq \min\{r, 20 - r\}$;
- (2) $r + a \equiv 0 \pmod{2}$; if $\delta_\varphi = 0$, then $r \equiv 2 \pmod{4}$;
- (3) if $a = 0$, then $(\delta_\varphi = 0, r \equiv 2 \pmod{8})$;
- (4) if $a = 1$, then $r \equiv 1, 3 \pmod{8}$;
- (5) if $(a = 2, r \equiv 6 \pmod{8})$, then $\delta_\varphi = 0$;
- (6) if $(a = r, \delta_\varphi = 0)$, then $r \equiv 2 \pmod{8}$;
- (7) if $(a = 20 - r, \delta_\varphi = 0)$, then $r \equiv 2 \pmod{8}$.

I. Conditions on $\delta_F, \delta_{\varphi F}$:

General conditions:

- (1) if $\delta_F = 0$, then $\delta_\varphi = 1$;
- (2) if $\delta_{\varphi F} = 0$, then $(\delta_F = 0, \delta_\varphi = 1, r \equiv 1 \pmod{4})$.

Relations near the boundary $a = 20 - r$:

- (3) if $a = 20 - r$, then $\delta_F = 0$;
- (4) if $(a = 20 - r, \delta_{\varphi F} = 0)$, then $r \equiv 1 \pmod{8}$.

Relations near the boundary $a = 0$:

- (5) if $a = 0$, then $\delta_F = 1$;
- (6) if $(a = 1, \delta_F = 0)$, then $(\delta_{\varphi F} = 0, r \equiv 1 \pmod{8})$;
- (7) if $(a = 2, \delta_F = 0)$, then $r \equiv 0, 2 \pmod{8}$;
- (8) if $(a = 3, \delta_F = 0, r \equiv 5 \pmod{8})$, then $\delta_{\varphi F} = 0$.

It is easy to enumerate all invariants (2.4) satisfying these conditions. They are all presented in Figure 1 for $\delta_F = 0$, and in Figure 2 for $\delta_F = 1$. Thus Figures 1 and 2 enumerate all connected components of moduli.

In [19], instead of δ_F, δ_φ and $\delta_{\varphi F}$, other invariants were also used. They are: the subgroup $H \subset (\mathbb{Z}/2\mathbb{Z})\text{cl}(F) = (2S^*/2S)$, the invariant $\delta_{\varphi S} \in \{0, 1\}$, the characteristic element $v \in H$ when $\delta_{\varphi S} = 0$ (the v is not defined when $\delta_{\varphi S} = 1$). They are related in the obvious way:




FIGURE 1. $\mathbb{F}_3$: The complete list of invariants 2.4 of connected components of moduli with $\delta_F = 0$ (then $\delta_\varphi = 1$).




FIGURE 2. $\mathbb{F}_3$: The complete list of invariants 2.4 of connected components of moduli with $\delta_F = 1$ (then $\delta_{\varphi F} = 1$).

$H = 0$ iff $\delta_F = 1$; $\delta_{\varphi S} = 0$ and $v = 0 \pmod{2S}$ iff $\delta_\varphi = 0$; $\delta_{\varphi S} = 0$ and $v \sim F \pmod{2S}$ iff $\delta_{\varphi F} = 0$; otherwise (if $\delta_\varphi = \delta_{\varphi F} = 1$) the invariant $\delta_{\varphi S} = 1$.

By simple direct calculations or applying ([19], Theorem 11 and Corollary 13), one finds how invariants (2.4) of related involutions φ and $\tilde{\varphi} = \tau\varphi$ can be calculated from each other:

$$(2.5) \quad \begin{aligned} r(\varphi) + r(\tilde{\varphi}) &= 19; \quad a(\varphi) + \delta_F(\varphi) = a(\tilde{\varphi}) + \delta_F(\tilde{\varphi}); \\ \delta_F(\varphi) + \delta_F(\tilde{\varphi}) &= 1; \quad \delta_{\tilde{\varphi}} = \delta_{\varphi F}; \quad \delta_{\tilde{\varphi} F} = \delta_\varphi. \end{aligned}$$

One can clearly see all these relations looking at Figures 1, 2.

Thus, we finally obtain the main result which enumerates connected components of moduli we are looking for.

Theorem 1. *There are exactly 102 connected components of moduli of positive real non-singular curves $A \in |-2K_{\mathbb{F}_3}|$ up to the action of $\text{Aut}(\mathbb{F}_3/\mathbb{R})$, equivalently of real K3 surfaces with non-symplectic involution of type $(3, 1, 1)$, equivalently of real elliptic K3 surfaces with section and no reducible fibres except one of the type $\tilde{\mathbb{A}}_1$ or $\tilde{\mathbb{A}}_2$. They are labelled by their natural invariants (2.4) satisfying relations 0.1–7 and I.1–8 above enumerated in Figures 1 and 2.*

Identifying related pairs of positive curves and anti-holomorphic involutions of K3, there are exactly 51 connected components of moduli of real non-singular curves $A \in |-2K_{\mathbb{F}_3}|$ up to the action of $\text{Aut}(\mathbb{F}_3/\mathbb{R})$, equivalently of related pairs (X, τ, φ) and $(X, \tau, \tilde{\varphi} = \tau\varphi)$ of real K3 surfaces with non-symplectic involution of type $(3, 1, 1)$, equivalently of related by the inverse map pairs of real elliptic K3 surfaces with section and no reducible fibres except one of the type $\tilde{\mathbb{A}}_1$ or $\tilde{\mathbb{A}}_2$. They are labelled by their related by (2.5) pairs of natural invariants (2.4) satisfying relations 0.1–7 and I.1–8 above enumerated in Figures 1 and 2.

Below we consider geometric interpretation of the invariants (2.4) in terms of K3 and curves on $\mathbb{F}_3$.

2.3. Geometric interpretation of the invariants (2.4) in terms of K3. We have the following geometric interpretation of the invariants $(r, a, \delta_\varphi, \delta_{\varphi F})$ in terms of the real K3 surface (X, φ) (see [15] and also [12]). We denote by T_g an orientable closed surface of genus g . We have

$$(2.6) \quad X_\varphi(\mathbb{R}) = \begin{cases} \emptyset & \text{if } (r, a, \delta_\varphi) = (10, 10, 0) \\ T_1 \amalg T_1 & \text{if } (r, a, \delta_\varphi) = (10, 8, 0) \\ T_g \amalg kT_0 & \text{otherwise,} \end{cases}$$

where

$$g = \frac{22 - r - a}{2}, \quad k = \frac{r - a}{2};$$

$$(2.7) \quad X_\varphi(\mathbb{R}) \sim 0 \pmod{2} \quad \text{in } H_2(X(\mathbb{C}); \mathbb{Z}),$$

if and only if $\delta_\varphi = 0$, and

$$(2.8) \quad X_\varphi(\mathbb{R}) \sim F(\mathbb{C}) \pmod{2} \quad \text{in } H_2(X(\mathbb{C}); \mathbb{Z}),$$

if and only if $\delta_{\varphi F} = 0$.

For the invariant δ_F , we have the following theorem (see Theorem 16 in [19]).

Theorem 2 ([19]). *Let X be a compact Kählerian surface with an anti-holomorphic involution φ , and $X_\varphi(\mathbb{R}) = X(\mathbb{C})^\varphi$ the set of real points. We assume that $H_1(X(\mathbb{C}); \mathbb{Z}) = 0$ and $X_\varphi(\mathbb{R}) \neq \emptyset$. Let C be a 1-dimensional complex submanifold of X with $\varphi(C) = C$, and $C(\mathbb{R})$ be the fixed point set of φ on C . Let $cl(C) \in H_2(X(\mathbb{C}); \mathbb{Z})$ and $cl(C(\mathbb{R})) \in H_1(X_\varphi(\mathbb{R}); \mathbb{Z}/2\mathbb{Z})$ denote the homology classes represented by C and $C(\mathbb{R})$ respectively.*

Then, $cl(C) \cdot H_2(X(\mathbb{C}); \mathbb{Z})_\varphi \equiv 0 \pmod{2}$ (equivalently, $\delta_C(\varphi) = 0$) if and only if $cl(C(\mathbb{R})) = 0$ in $H_1(X_\varphi(\mathbb{R}); \mathbb{Z}/2\mathbb{Z})$ (as usual, $H_2(X(\mathbb{C}); \mathbb{Z})_\varphi$ denotes the eigenvalue -1 part of φ).

Thus, if F is irreducible, then $\delta_F = 0$ if and only if $F(\mathbb{R}) = 0$ in $H_1(X_\varphi(\mathbb{R}); \mathbb{Z}/2\mathbb{Z})$. Changing the real structure a little (in the same connected component of moduli), we can always make F irreducible, $F \cong \mathbb{P}^1$, and it is then a complex submanifold of $X(\mathbb{C})$. By deformation argument, we then have the same statement for a reducible F . Since $X_\varphi(\mathbb{R})$ is always non-empty, we get at any case

$$(2.9) \quad \delta_F = 0 \text{ if and only if } cl(F(\mathbb{R})) = 0 \text{ in } H_1(X_\varphi(\mathbb{R}); \mathbb{Z}/2\mathbb{Z}).$$

Thus, all invariants (2.4) have very clear geometric meaning in terms of the real K3 surface (X, φ) .

Theorem 3. *The connected component of moduli of a real K3 surface with non-symplectic involution (X, τ, φ) of the type $(3, 1, 1)$ (equivalently of a positive real non-singular curve $A \in |-2K_{\mathbb{F}_3}|$) is determined by the geometric invariants: topology of the real part $X_\varphi(\mathbb{R})$, if $F_\varphi(\mathbb{R}) \sim 0 \pmod{2}$ in $H_1(X_\varphi(\mathbb{R}); \mathbb{Z})$, if $X_\varphi(\mathbb{R}) \sim 0 \pmod{2}$ in $H_2(X(\mathbb{C}); \mathbb{Z})$, if $X_\varphi(\mathbb{R}) \sim F(\mathbb{C}) \pmod{2}$ in $H_2(X(\mathbb{C}); \mathbb{Z})$.*

See (2.6), (2.7), (2.8) and (2.9) about their exact relation with the complete invariants (2.4) of the connected component of moduli.

Taking the related involution $\tilde{\varphi} = \tau\varphi$ instead of φ , we obtain geometric interpretation of the invariants (2.4) in terms of the real K3 surface $(X, \tilde{\varphi})$.

Theorem 4. *The connected component of moduli of a real K3 surface with non-symplectic involution (X, τ, φ) of type $(3, 1, 1)$ (equivalently of a positive real non-singular curve $A \in |-2K_{\mathbb{F}_3}|$) is determined by the geometric invariants: topology of the real part $X_{\tilde{\varphi}}(\mathbb{R})$, if $F_{\tilde{\varphi}}(\mathbb{R}) \sim 0 \pmod{2}$ in $H_1(X_{\tilde{\varphi}}(\mathbb{R}); \mathbb{Z})$, if $X_{\tilde{\varphi}}(\mathbb{R}) \sim 0 \pmod{2}$ in $H_2(X(\mathbb{C}); \mathbb{Z})$, if $X_{\tilde{\varphi}}(\mathbb{R}) \sim F(\mathbb{C}) \pmod{2}$ in $H_2(X(\mathbb{C}); \mathbb{Z})$.*

See (2.6), (2.7), (2.8) (2.9) and (2.5) about their exact relation with the complete invariants (2.4) of the connected component of moduli.

2.4. Geometric interpretation of the invariants (2.4) in terms of $A \in |-2K_{\mathbb{F}_3}|$. Next we consider the dividedness of the fixed point curve of a real K3 surface with a non-symplectic involution.

Let A be a possibly disconnected non-singular complex compact curve with an anti-holomorphic involution φ , and $A(\mathbb{R})$ the fixed point set of φ . We remind that (A, φ) (or simply, A) is called *dividing* if $A(\mathbb{R}) = 0 \in H_1(A; \mathbb{Z}/2\mathbb{Z})$. We have the following result which had been partially used in [19] (see the proof of Theorem 26 in [19]).

Theorem 5. *Let (X, τ, φ) be a real K3 surface with non-symplectic involution of type $(S, \theta) = (H_2(X(\mathbb{C}); \mathbb{Z})^\tau, \varphi|_{H_2(X(\mathbb{C}); \mathbb{Z})^\tau})$ (see [19]); let $A = X^\tau$ be the fixed point set of τ in X . Assume that A is non-empty. Then, the real curve A is dividing if and only if $\delta_{\varphi S} = 0$ (equivalently, $A(\mathbb{R}) \pmod{2} \in H_2(X(\mathbb{C}); \mathbb{Z})^\tau \pmod{2}$).*

Proof. The main method is a so-called Donaldson's trick (see [5] or [4]). It is known that the K3 surface X is algebraic. Hence, we can take a hyperplane section class h in S . We denote $L = H_2(X(\mathbb{C}); \mathbb{Z})$. Since $\varphi_*(h) = -h$, h is contained in $S_- = L^\tau \cap L_\varphi$. Moreover, h is contained in the Kähler cone $\mathcal{C}_X^+$ of X . (See [3], p.242, for example.) Thus we see $S_- \cap \mathcal{C}_X^+ \neq \emptyset$. Let I be the complex structure of the K3 surface X and take a Kähler class c of (X, I) in $((L^\tau \cap L_\varphi) \otimes \mathbb{R}) \cap \mathcal{C}_X^+$. Since $c_1(X) = 0$, by the Calabi-Yau theorem (for example, [2], Theorem 11.15) 0 is the Ricci form of one and only one Kähler metric in the class c . Thus we can take the unique Kähler form P with zero Ricci form in the class c , where $c = [P]$. Let g be the Kähler metric whose Kähler form is P , i.e., $P = g(I(\cdot), \cdot)$. By the uniqueness of the Ricci flat Kähler form, we have $\tau^*P = P$ and $\varphi^*P = -P$. Since τ is holomorphic on (X, I) , we have $\tau^*g = g$. And, since φ is anti-holomorphic on (X, I) , we have $\varphi^*g = g$.

Now we can take a nowhere vanishing holomorphic 2-form ω_I on (X, I) such that $\tau^*\omega_I = -\omega_I$ and $\varphi^*(\omega_I) = \overline{\omega_I}$. We define the real 2-forms Q and R on X to be $\omega_I = Q + \sqrt{-1}R$. Then we have

$$\tau^*Q = -Q, \quad \tau^*R = -R, \quad \varphi^*Q = Q \quad \text{and} \quad \varphi^*R = -R.$$

Moreover, we may assume $2(P \wedge P) = \omega_I \wedge \overline{\omega_I}$, equivalently, $Q \wedge Q = P \wedge P$.

We define the new complex structures J and K on X by

$$Q = g(J(\cdot), \cdot) \text{ and } R = g(K(\cdot), \cdot).$$

(Then, g is a hyperKähler metric on X . See [9], Remark 2.4.) By the above we can verify that τ is anti-holomorphic and φ is holomorphic for (X, J) . We see $\omega_J := R + \sqrt{-1}P$ is a nowhere vanishing holomorphic 2-form on (X, J) . We have $\varphi^*(\omega_J) = -\omega_J$, i.e., φ is non-symplectic on the K3 surface (X, J) . We get a new real K3 surface with a non-symplectic involution $((X, J), \varphi, \tau)$.

Then the fixed point set $X_\varphi(\mathbb{R})$ of φ is a complex 1-dimensional submanifold of (X, J) ([16], p.1424). The fixed point set of τ in $X_\varphi(\mathbb{R})$ is $A(\mathbb{R})$. Thus, we can apply Theorem 2 to the K3 surface (X, J) , the anti-holomorphic involution τ and the complex curve $C := X_\varphi(\mathbb{R})$. Thus we conclude that the following two conditions are equivalent.

- (1) $cl(X_\varphi(\mathbb{R})) \cdot x \equiv 0 \pmod{2} \forall x \in L_\tau$.
- (2) $cl(A(\mathbb{R})) = 0 \in H_1(A; \mathbb{Z}/2\mathbb{Z})$.

The condition (1) is equivalent to the condition $\delta_{\varphi S} = 0$. This completes the theorem. $\square$

In our case, $A = A_0 \amalg A_1$ where $A_0 \cong \mathbb{P}^1$ and A_1 has genus 9. Thus, A_0 is always dividing, and A is dividing if and only if A_1 is dividing which makes sense also for $A = s + A_1 \in |-2K_{\mathbb{F}_3}|$. Thus, applying Theorem 5, we get in our case that

$$(2.10) \quad A_1 \text{ is dividing if and only if either } (\delta_F = 1, \delta_\varphi = 0) \text{ or } \delta_F = \delta_{\varphi F} = 0.$$

Let us consider real $\mathbb{F}_3$. Since $s^2 = -3$ is odd, it follows that $s(\mathbb{R})^2 \equiv 1 \pmod{2}$ in $H_1(\mathbb{F}_3(\mathbb{R}); \mathbb{Z}/2\mathbb{Z})$. It follows that $\mathbb{F}_3(\mathbb{R})$ is non-empty, and it is a non-orientable surface. Rational pencil of $\mathbb{F}_3$ gives then a pencil on $\mathbb{F}_3(\mathbb{R})$ by circles over the circle $s(\mathbb{R})$. Thus, $\mathbb{F}_3(\mathbb{R})$ is Klein bottle. Further we denote by c a real (i. e. over a point of $s(\mathbb{R})$) fibre of the pencil. Then $c(\mathbb{R})$ is a circle. The surface Y (the blow-up of $\mathbb{F}_3$ at $A_1 \cap s$) has $Y(\mathbb{R})$ which is a non-orientable surface of genus 2 ($\chi(Y(\mathbb{R})) = -1$).

We consider isotopy type of the curve $A(\mathbb{R}) = s(\mathbb{R}) \cup A_1(\mathbb{R})$ in $\mathbb{F}_3(\mathbb{R})$. By an *oval*, we mean a connected component of $A_1(\mathbb{R})$ which bounds a disk in $\mathbb{F}_3(\mathbb{R})$. Each otherwise connected component is called a *non-oval*.

We know that $A_1 \cdot s = 1$ and $A_1 \cdot c = 3$. Moreover, we know the geometric interpretation of invariants (2.4) in terms of the double covering K3 from Sect. 2.3. Using this information, by elementary 2-dimensional topology considerations, we obtained all possible pictures.

Let o_1 be the connected component (a circle) of $A_1(\mathbb{R})$ which contains the unique point $A_1 \cap s$. The o_1 is a non-oval, and there are two cases.

Case 1. $A_1(\mathbb{R})$ consists of the non-oval o_1 and $g - 1 + k$ ovals. In case of $\delta_F = 0$, the positive curve A^+ is shown as shadow in Figure 3. As usual, $g = (22 - r - a)/2$ and $k = (r - a)/2$. This case does not include invariants $(r, a, \delta_{\varphi F}) = (9, 9, 0)$ which will appear in the case 2 below. The negative part $A^- = \overline{\mathbb{F}_3(\mathbb{R})} - A^+$ gives then the positive curve for the related connected component of moduli with $\delta_F = 1$. By (2.5), one can write down the corresponding related invariants (2.4) of this positive curve.

In this case, the blow up $\pi(X_\varphi(\mathbb{R}))$ of A^+ consists of a closed cylinder with $g - 1$ holes, and k closed disks. Two boundary circles of the cylinder are $s(\mathbb{R})$ and o_1 . The blow up $\pi(X_{\tilde{\varphi}}(\mathbb{R}))$ of A^- consists of a real projective plane with $k + 2$ holes, and $g - 1$ disks. The boundaries of




FIGURE 3. The positive curve A^+ for $\delta_F = 0$ and $(r, a, \delta_{\varphi F}) \neq (9, 9, 0)$ (see Figure 1), here $g = (22 - r - a)/2$, $k = (r - a)/2$.

two these holes are $s(\mathbb{R})$ and o_1 . Identifying the boundaries of the corresponding disks and holes of $\pi(X_\varphi(\mathbb{R}))$ and $\pi(X_{\bar{\varphi}}(\mathbb{R}))$, one obtains $Y(\mathbb{R})$.

Case 2. $A_1(\mathbb{R})$ consists of the non-oval o_1 and another non-oval o_2 . In case of $\delta_F = 0$, the positive curve A^+ is shown as shadow in Figure 4. In this case, $(r, a, \delta_{\varphi F}) = (9, 9, 0)$. The negative part $A^- = \overline{\mathbb{F}_3(\mathbb{R})} - A^+$ then corresponds to the positive curve with the invariants $\delta_F = 1$ and $(r, a, \delta_{\varphi}) = (10, 8, 0)$.

In this case, the blow up $\pi(X_\varphi(\mathbb{R}))$ of A^+ is a closed disk with 2 holes. The boundary circles of the holes of the disk are $s(\mathbb{R})$ and o_1 . The boundary of the disk is o_2 . The blow up $\pi(X_{\bar{\varphi}}(\mathbb{R}))$ of A^- consists of a closed cylinder and a Möbius band. Two boundary circles of the cylinder are $s(\mathbb{R})$ and o_1 . The boundary of the Möbius band is o_2 .

These pictures and (2.10) show that the isotopy type of A^+ together with dividedness determine the invariants (2.4). By Theorem 1 we finally get

Theorem 6. *The connected component of moduli of a positive curve A^+ of a real non-singular curve $A = s + A_1 \in |-2K_{\mathbb{F}_3}|$ is determined by the isotopy type of A^+ and by dividedness by $A_1(\mathbb{R})$ of $A_1(\mathbb{C})$.*

All possibilities are presented in Figures 1 — 4 and (2.10).

We also have similar result in terms of the blow up surface Y (see its exact description above).

Theorem 7. *Let Y be the blow up of $\mathbb{F}_3$ at a point of the exceptional section of $\mathbb{F}_3$.*

The connected component of moduli of a positive curve A^+ of a real non-singular curve $A \in |-2K_Y|$ is determined by the topological type of A^+ and by dividedness by $A(\mathbb{R})$ of $A(\mathbb{C})$.

All possibilities are presented in Figures 1 — 4 and (2.10).




FIGURE 4. The positive curve A^+ for $\delta_F = 0$ and $(r, a, \delta_{\varphi_F}) = (9, 9, 0)$ (see Figure 1).

One can consider other important applications of our results.

Remark 8. *Let us take the contraction of the exceptional curve F of Y . Then one obtains $\mathbb{F}_4$. The image A' of A gives then a curve $A' \in |-2K_{\mathbb{F}_4}|$ which is non-singular except one quadratic (possibly degenerate) singular point which is the image of F .*

Thus, we obtain classification of connected components of moduli (up to $\text{Aut}(\mathbb{F}_4/\mathbb{R})$) of real curves $A' \in |-2K_{\mathbb{F}_4}|$ which are non-singular except one quadratic singular point (the blow up at this point resolves this singularity).

We leave details to the interested reader.

Remark 9. *Let us consider the contraction $\sigma : Y \rightarrow Z$ of the curve $A_0 \subset Y$ with $A_0^2 = -4$. The surface Z is a log del Pezzo surface of index ≤ 2 . Then image $\sigma(A_1)$ gives a non-singular curve in $|-2K_Z|$, and vice versa. See [1] for details. By classification of log del Pezzo surfaces of index ≤ 2 in [1], the surface Z is characterized by the properties: Z has no singular points except one log-terminal singular point of index two having the type K_1 (it is the image of A_0), moreover the genus $g(A_1) = 9$.*

Thus, we obtain classification of connected components of moduli (up to $\text{Aut}(Z/\mathbb{R})$) of real non-singular curves $A_1 \in |-2K_Z|$ where Z is a real del Pezzo surface having no singular points except one log-terminal singular point of index 2 of the type K_1 , and the genus $g(A_1) = 9$.

We leave details to the interested reader.

3. CONNECTED COMPONENTS OF MODULI OF REAL NON-SINGULAR CURVES IN $|-2K_{\mathbb{F}_2}|$

Here we consider the case of $\mathbb{F}_2$. We shall see that this case is more difficult, more interesting, very classical, and very rich.

3.1. Reduction to real K3 surfaces with degenerate non-symplectic involution of type $(2, 2, 0)$. Here we consider $Y = \mathbb{F}_2$. Like above, c denotes the fibre of the rational pencil of $\mathbb{F}_2$, and s its exceptional section. We have $c^2 = 0$, $s^2 = -2$ and $c \cdot s = 1$. We have $-2K_{\mathbb{F}_2} = 8c + 4s$.

Let $A \in |-2K_{\mathbb{F}_2}|$ be a non-singular curve. It is irreducible of genus 9. We have $A \cdot s = 0$, and A does not intersect s .

Let $\pi : X \rightarrow Y = \mathbb{F}_2$ be the double covering of Y ramified in $A \subset Y$. The X is a K3 surface with the non-symplectic involution τ of the double covering. Then $A = X^\tau$ is identified with the fixed points set of the involution, and $\mathbb{F}_2 = Y = X/\{1, \tau\}$ is the quotient space. We denote $C = \pi^{-1}(c)$, $F = \pi^{-1}(s)$. Since $s \cong \mathbb{P}^1$ and A does not intersect s , the curve $F = F_1 + F_2$ is disjoint union of two non-singular rational curves which are conjugate by τ . We have $F_1^2 = F_2^2 = -2$, $F_1 \cdot F_2 = 0$, $C \cdot F_1 = C \cdot F_2 = 1$. Here the numbering by 1 and 2 of the curves F_1 and F_2 is not canonical, of course. The Gram matrix of C , F_1 and F_2 has the determinant 4. Since there are no hyperbolic even unimodular lattices of the rank 3 (its signature must be 0 mod 8), it follows that C , F_1 and F_2 generate a primitive sublattice in the Picard lattice S_X of X .

Since Y has Picard number 2, it follows that C and $F = F_1 + F_2$ generate the eigenvalue 1 sublattice $S = S_X^\tau = H_2(X(\mathbb{C}); \mathbb{Z})^\tau = \mathbb{Z}cl(C) + \mathbb{Z}cl(F)$ of τ . We have $r(S) = \text{rk } S = 2$, $S^*/S \cong (\mathbb{Z}/2\mathbb{Z})^2$ is a 2-elementary group of the rank $a(S) = 2$, we have $(x^*)^2 \in \mathbb{Z}$ for any $x^* \in S^* = S/2$. It follows that the invariant $\delta(S) = 0$. Thus, the main invariants of τ are $(r(S), a(S), \delta(S)) = (2, 2, 0)$. Moreover, in our case, all exceptional curves on X such that their image by the quotient map $\pi : X \rightarrow Y$ gives exceptional curves on $Y = \mathbb{F}_2$, are the non-singular rational curves F_1 and F_2 on X such that $F_1^2 = F_2^2 = -2$, $F_1 \cdot F_2 = 0$ and $\tau(F_1) = F_2$. Other speaking, all pre-images by π of exceptional curves on Y consist of the exceptional curves F_1 and F_2 .

Vice versa, let us consider a K3 surface with non-symplectic involution (X, τ) where $S = (S_X)^\tau$ is a 2-elementary lattice with the same invariants $(r(S), a(S), \delta(S)) = (2, 2, 0)$. By simple direct considerations, or by using general classification results of [1], there are two cases for X and the quotient map $\pi : X \rightarrow Y = X/\{1, \tau\}$. In both cases $Y = X/\{1, \tau\}$ is a rational surface with Picard number $r(S) = 2$, and $A = X^\tau$ gives a non-singular irreducible curve $A = \pi(A) \in |-2K_Y|$ of genus 9. Thus, π is a double covering ramified in a non-singular curve $A \in |-2K_Y|$. In first case, Y has no exceptional curves. In second case, Y has exactly one exceptional curve s with $s^2 = -2$. By classification of surfaces (e. g. see [23]), then $Y \cong \mathbb{F}_0 = \mathbb{P}^1 \times \mathbb{P}^1$ in first case, and $Y = \mathbb{F}_2$ in second case. First case is called *non-degenerate*. Second case is called *degenerate*.

The first, non-degenerate case, had been studied over $\mathbb{R}$ in [19].

The second, degenerate case, we consider here. We see that non-singular curves $A \in |-2K_{\mathbb{F}_2}|$ and K3 surfaces with degenerate non-symplectic involution (X, τ) of type $(2, 2, 0)$ are in one-to-one correspondence. They have the same moduli and their connected components.

Now let us assume that $Y = \mathbb{F}_2$ and a non-singular curve $A \in |-2K_Y|$ are real. Let θ be the corresponding anti-holomorphic involution on Y . Since Y has the unique rational pencil, $|c|$, and the unique exceptional curve, s , then $\theta(cl(c)) = -cl(c)$, $\theta(cl(s)) = -cl(s)$, i. e. θ acts as (-1) in $H_2(Y(\mathbb{C}); \mathbb{Z})$. Let φ be the lift of θ on X . The φ is defined uniquely up to replacement by the related $\tilde{\varphi} = \tau\varphi$. Replacing φ by $\tilde{\varphi}$ if necessary, we can always assume

that $\varphi(F_1) = F_1$ and $\varphi(F_2) = F_2$. It follows that φ acts as (-1) in the part of $H_2(X(\mathbb{C}); \mathbb{Z})$ generated by C , F_1 and F_2 . Thus the action of φ in $H_2(X(\mathbb{C}); \mathbb{C})$ has at least 3 linearly independent eigenvectors with the eigenvalue (-1) . If φ has order 4, then $\varphi^2 = \tau$, and φ must have twenty linearly independent eigenvectors with eigenvalues $\pm\sqrt{-1}$ in the eigenvalue -1 part $H_2(X(\mathbb{C}); \mathbb{C})_\tau$ of the action of τ . Thus, we obtain more than $\text{rk } H_2(X(\mathbb{C}); \mathbb{Z})$ linearly independent eigenvectors for the action of φ which is impossible. This shows that φ has order 2. The involutions φ and τ obviously commute.

Thus, there exists a unique lift φ of θ to an anti-holomorphic involution φ of X which is determined by the condition $\varphi(F_1) = F_1$ and $\varphi(F_2) = F_2$. Then φ acts as -1 in the $\mathbb{Z}$ -submodule $[C, F_1, F_2]$ of $H_2(X(\mathbb{C}); \mathbb{Z})$ generated by C , F_1 and F_2 . For the related involution $\tilde{\varphi} = \tau\varphi$ we have $\tilde{\varphi}(F_1) = F_2$ and $\tilde{\varphi}(F_2) = F_1$, $\tilde{\varphi}$ acts as (-1) in $H_2(X(\mathbb{C}); \mathbb{Z})^\tau$, and $\tilde{\varphi}(cl(F_1) - cl(F_2)) = cl(F_1) - cl(F_2)$ where $cl(F_1) - cl(F_2) \in H_2(X(\mathbb{C}); \mathbb{Z})_\tau$.

Thus, classifying connected components of moduli of real K3 surfaces with degenerate non-symplectic involution (X, τ, φ) of type $(2, 2, 0)$, we actually solve more delicate problem of classification of connected components of moduli of positive real non-singular curves A^+ where $A \in |-2K_{\mathbb{F}_2}|$ is non-singular. Here A^+ is determined by a choice of one of two related anti-holomorphic involutions φ or $\tilde{\varphi}$ of the double covering X . If $Y(\mathbb{R}) \neq \emptyset$, we can identify A^+ with the closure of one of two parts of $Y(\mathbb{R}) - A(\mathbb{R})$ which is $A^+ = \pi(X_\varphi(\mathbb{R}))$. Then $A^- = \overline{Y(\mathbb{R}) - A^+}$ is $A^- = \pi(X_{\tilde{\varphi}}(\mathbb{R}))$.

If $Y(\mathbb{R}) \neq \emptyset$, then $s(\mathbb{R}) \neq \emptyset$, and the part A^+ corresponding to the choice of φ above is determined by the condition $s(\mathbb{R}) \subset A^+$. Further we always assume that the positive curve A^+ is chosen in this way: it corresponds to φ which acts on $[C, F_1, F_2]$ as (-1) .

3.2. Enumeration of connected components of moduli of real K3 surfaces with degenerate non-symplectic involution of type $(2, 2, 0)$. We correspond to a real K3 surface with non-symplectic involution (X, τ, φ) (or $(X, \tau, \tilde{\varphi})$) of type $(2, 2, 0)$ the isomorphism class of the action (L, τ, φ) of the involutions τ and φ on the homology lattice $L = H_2(X(\mathbb{C}); \mathbb{Z})$ (which is an even unimodular lattice of signature $(3, 19)$) where τ has type $(2, 2, 0)$ and φ has hyperbolic L^φ . Here τ and φ must act on $[C, F_1, F_2]$ as has been described above: $\tau(cl(C)) = cl(C)$, $\tau(cl(F_1)) = cl(F_2)$, $\tau(cl(F_2)) = cl(F_1)$ and φ is (-1) on $[C, F_1, F_2]$. Respectively $\tilde{\varphi} = \tau\varphi$ acts on $[C, F_1, F_2]$ as follows: $\tilde{\varphi}(cl(C)) = -cl(C)$, $\tilde{\varphi}(cl(F_1)) = -cl(F_2)$, $\tilde{\varphi}(cl(F_2)) = -cl(F_1)$. Any action with these properties takes place for some (X, τ, φ) (or $(X, \tau, \tilde{\varphi})$).

Two such actions (L, τ, φ) and (L', τ', φ') are called isomorphic if there exists an isomorphism $f : L \rightarrow L'$ of lattices such that $f\tau = \tau'f$, $f\varphi = \varphi'f$, $f(cl(C)) = cl(C)$, $f(cl(F_1)) = cl(F_1)$ and $f(cl(F_2)) = cl(F_2)$ (i. e. f is identical on $[C, F_1, F_2]$). Replacing f by $f\tau$, we can always more generally assume that $f(\{cl(F_1), cl(F_2)\}) = \{cl(F_1), cl(F_2)\}$.

Since our case is degenerate (because Y has real exceptional curves with square (-2)), we cannot apply results of ([19], Sect. 2.1) where only non-degenerate case had been considered. In spite of that, in our very special case similar result is valid.

Theorem 10. *The connected component of moduli of a real K3 surface (X, τ, φ) (or $(X, \tau, \tilde{\varphi} = \tau\varphi)$) with degenerate non-symplectic involution of type $(2, 2, 0)$ is determined by the isomorphism class of the action of τ and φ (or $\tilde{\varphi}$) on the homology lattice $L = H_2(X; \mathbb{Z})$.*

Proof. We have $L^\tau = \mathbb{Z}cl(C) + \mathbb{Z}(cl(F_1) + cl(F_2))$. Its orthogonal complement in $[C, F_1, F_2]$ is $\mathbb{Z}f$ where $f = cl(F_1) - cl(F_2)$ and $f^2 = -4$. We denote by T the orthogonal complement to $[C, F_1, F_2]$ in L .

Assume that $\delta \in L$ satisfies the properties: $\delta \perp L^\tau$, the sublattice in L generated by $[C, F_1, F_2]$ and δ is hyperbolic, and $\delta^2 = -2$.

Then $\delta = \alpha f + \beta \delta_2$ where α, β are rational numbers, $\delta_2 \in T$ is primitive and $\delta_2^2 < 0$.

We claim that then either $\alpha = 0$ and $\delta = \pm \delta_2 \in T$, or $\alpha = \pm(1/2)$, $\beta = \pm(1/2)$ and $f^2 = \delta_2^2 = -4$.

Really, if $\alpha = 0$, then $\delta = \beta \delta_2 \in T$ is primitive, and then $\beta = \pm 1$.

Assume that $\alpha \neq 0$. We have $F_1 \cdot \delta = -2\alpha \in \mathbb{Z}$, and $\alpha = a/2$ where $a \in \mathbb{Z}$ and $a \neq 0$. Then $\delta = (a/2)f + \beta \delta_2$ and $2\delta - af = 2\beta \delta_2 \in L$. It follows that $2\beta \in \mathbb{Z}$ and $\beta = b/2$ where $b \in \mathbb{Z}$. Thus $\delta = (a/2)f + (b/2)\delta_2$ where a and b are non-zero integers. We have $\delta^2 = -2 = -a^2 + (b^2/4)\delta_2^2$ where $\delta_2^2 < 0$ is even. It follows that $\delta_2^2 = -4$ and $a = \pm 1$, $b = \pm 1$.

Thus, $\delta = (\pm f \pm \delta_2)/2$ where $f \in [C, F_1, F_2]$ and $\delta_2 \in T$ are elements with square -4 . We have $[C, F_1, F_2] \cdot \delta = [C, F_1, F_2] \cdot f/2 \in \mathbb{Z}$ and $T \cdot \delta = T \cdot \delta_2/2 \in \mathbb{Z}$. Thus, $[C, F_1, F_2] \cdot f \equiv 0 \pmod{2}$ and $T \cdot \delta_2 \equiv 0 \pmod{2}$. Thus, $f' = \pm f$ is a root of $[C, E_1, E_2]$ with square (-4) , $\delta'_2 = \pm \delta_2$ is a root of T with square (-4) , and $\delta = (f' + \delta'_2)/2$ is the half sum of these roots. Moreover, $\delta' = (f' - \delta'_2)/2 \in L$ and $(\delta')^2 = -2$.

If the mirror $\delta^\perp$ of δ divides the corresponding period domain of (X, τ, φ) of our type in two disjoint half-spaces, they will be conjugate by the reflection in δ in the first case, and by the composition of reflections in δ and δ' in the second case. It is easy to see that these automorphisms will conjugate periods of isomorphic K3 surfaces with non-symplectic involution.

The remaining considerations are the same as for the non-degenerate case in the proof of Theorem 1 in [19].

□

Remark 11. *Similar (properly formulated) result can be proved for real K3 surfaces with degenerate non-symplectic involution and the root invariant (K, ξ) where K is a 2-elementary root lattice (i. e. it is orthogonal sum of A_1, D_{2k}, E_7, E_8). See [1] for terminology and classification results. For our $\mathbb{F}_2$ case, K is A_1 . For non-degenerate case, K is zero.*

Unfortunately, it seems, similar to Theorem 10 result holds only for very special classes of K3 surfaces with degenerate non-symplectic involution. It would be interesting to find all these nice cases.

Thus, connected components of moduli of positive real non-singular curves $A \in |-2K_Y|$ (further always $Y = \mathbb{F}_2$) and, equivalently, of real K3 surfaces with degenerate non-symplectic involution (X, τ, φ) of type $(2, 2, 0)$ with our choice of φ between two related anti-holomorphic involutions are in one-to-one correspondence with isomorphism classes of triplets

$$(3.1) \quad (L, \tau, \varphi)$$

described above.

They can be enumerated using general results of [17] about classification of integral involutions with conditions. On the other hand, this case (like $\mathbb{F}_3$ above) can be also reduced to classification of integral polarized involutions (see Sect. 3 in [15]) which is much simpler

since all necessary calculations are already done in ([15], Theorem 3.4.3). We prefer the second possibility.

We can decompose $[C, F_1, F_2] = [C, F_1] \oplus [\tilde{F}_1]$ where $\tilde{F}_1 = 2C + (F_1 - F_2)$. Here $[C, F_1] = U$ is even unimodular lattice of signature $(1, 1)$ and $[\tilde{F}_1] = \langle -4 \rangle$ since $(2C + F_1 - F_2)^2 = -4$

A triplet (L, τ, φ) from (3.1) defines an integral polarized involution

$$(3.2) \quad (L_1, \varphi_1, cl(\tilde{F}_1)).$$

Here L_1 is the orthogonal complement to U in L (thus, it is an even unimodular lattice of signature $(2, 18)$, $\varphi_1 = \varphi|_{L_1}$ and $L_1^{\varphi_1}$ is hyperbolic, moreover, $\varphi_1(cl(\tilde{F}_1)) = -cl(\tilde{F}_1)$ and $\tilde{F}_1^2 = -4$.

Taking orthogonal sum of (3.2) with $U = [C, F_1]$, one obtains the triplet (3.1) where τ acts as $\tau(cl(C)) = cl(C)$, $\tau(cl(F_1)) = cl(F_2)$, $\tau(cl(F_2)) = cl(F_1)$, and τ is (-1) on $\tilde{F}_1^\perp$ in L_1 . Thus, their isomorphism classes are in one to one correspondence.

Genus invariants of integral polarized involutions (3.2) (and then (3.1)) were completely classified in Theorem 3.4.3 of [15]. Below we present these results. (One should multiply by (-1) the pairing in L_1 and apply Theorem 3.4.3 of [15] to $l_{(+)} = 18$, $l_{(-)} = 2$, $t_{(-)} = 1$, $t_{(+)} = r - 1$ and $n = 4$.)

The complete genus invariants of (3.2) and then (3.1) are

$$(3.3) \quad (r, a, \delta_\varphi; \delta_F, \delta_{\varphi F}).$$

Here $r = \text{rk } L^\varphi \in \mathbb{N}$; $((L^\varphi)^*/L^\varphi) = (\mathbb{Z}/2\mathbb{Z})^a$ where $a \geq 0$ is an integer; $\delta_\varphi \in \{0, 1\}$ is equal to 0 if and only if $x \cdot \varphi(x) \equiv 0 \pmod{2}$ for any $x \in L$. They are all invariants of the corresponding pair (L, φ) . Here $\delta_F \in \{0, 1\}$ is equal to 0 if and only if $F \cdot L_\varphi \equiv 0 \pmod{2}$; here $\delta_{\varphi F} \in \{0, 1\}$ is equal to 0 if and only if $x \cdot \varphi(x) \equiv x \cdot F$ for any $x \in L$. Here and in what follows, we always denote by L^φ and L_φ the eigenvalue 1 and -1 parts respectively of the action of an involution φ on a module L .

Strictly speaking, for invariants of integral polarized involutions (3.2), one should use $\tilde{F}_1$ instead of F , φ_1 instead of φ , and L_1 instead of L . But, $\tilde{F}_1 \equiv F = F_1 + F_2 \pmod{2}$ in L , and the invariants are the same: $\delta_F = \delta_{\tilde{F}_1}$, $\delta_\varphi = \delta_{\varphi_1}$, and $\delta_{\varphi F} = \delta_{\varphi_1 \tilde{F}_1}$.

The invariants (3.3) must satisfy the following relations which are sufficient and necessary for existence of a triplet (3.2) or the corresponding triplet (3.1) with these invariants.

0. Conditions on (r, a, δ_φ) :

- (1) $1 \leq r \leq 18$, $0 \leq a \leq \min\{r, 20 - r\}$;
- (2) $r + a \equiv 0 \pmod{2}$; if $\delta_\varphi = 0$, then $r \equiv 2 \pmod{4}$;
- (3) if $a = 0$, then $(\delta_\varphi = 0, r \equiv 2 \pmod{8})$;
- (4) if $a = 1$, then $r \equiv 1, 3 \pmod{8}$;
- (5) if $(a = 2, r \equiv 6 \pmod{8})$, then $\delta_\varphi = 0$;
- (6) if $(a = r, \delta_\varphi = 0)$, then $r \equiv 2 \pmod{8}$;
- (7) if $(a = 20 - r, \delta_\varphi = 0)$, then $r \equiv 2 \pmod{8}$.

I. Conditions on δ_F and $\delta_{\varphi F}$:

General conditions:

- (1) if $\delta_{\varphi F} = 0$, then $(\delta_F = 0, \delta_\varphi = 1, r \equiv 0 \pmod{4})$.

Relations near the boundary $a = 20 - r$:




FIGURE 5. $\mathbb{F}_2$: The complete list of the invariants 3.3 (i. e. in terms of φ) with $\delta_F = 0$ of the connected components of moduli.




FIGURE 6. $\mathbb{F}_2$: The complete list of the invariants 3.3 (i. e. in terms of φ) with $\delta_F = 1$ (then $\delta_{\varphi F} = 1$) of the connected components of moduli.

- (2) if $a = 20 - r$, then $\delta_F = 0$;
- (3) if $(a = 20 - r, \delta_{\varphi F} = 0)$, then $r \equiv 0 \pmod{8}$;
- (4) if $(a = 18 - r, \delta_F = 1, \delta_\varphi = 0)$, then $r \equiv 2 \pmod{8}$.

Relations near the boundary $a = 0$:

- (5) if $a = 0$, then $\delta_F = 1$;
- (6) if $a = 1$, then $\delta_F = 1$;
- (7) if $(a = 2, \delta_F = 0, r \equiv 2 \pmod{8})$, then $\delta_\varphi = 0$;
- (8) if $(a = 2, \delta_F = 0, r \equiv 0 \pmod{4})$, then $\delta_{\varphi F} = 0$.

It is easy to enumerate all invariants (3.3) satisfying these conditions. They are all presented in Figure 5 for $\delta_F = 0$, and in Figure 6 for $\delta_F = 1$.

Additional considerations show that the genus invariants (3.3) define the isomorphism class.

Lemma 12. *The genus invariants (3.3) determine the isomorphism class of (3.2) and of (3.1). Thus, they determine the connected component of moduli.*

Proof. If $20 - r \geq 4 + \delta_F$, the statement follows from Theorem 3.3.1 in [15].

Let us assume that $20 - r < 4 + \delta_F$. Equivalently, $r \geq 17 - \delta_F$. Let us denote $h = cl(\tilde{F}_1)$. We have $h^2 = -4$.

In all cases it is enough to prove (see [15] or Remark 1.6.2 in [17]) that the lattice $K = (L_1)_{\varphi_1, F}$ which is the orthogonal complement to h in $(L_1)_{\varphi_1}$ has the properties:

$$(3.4) \quad \begin{aligned} &K \text{ is unique in its genus and the canonical homomorphism} \\ &O(K) \rightarrow O(q_K) \text{ is epimorphic.} \end{aligned}$$

Here q_K denotes the discriminant quadratic form of K .

Further we denote by E_8 and E_7 the negative definite lattices defined by the Dynkin diagrams $\mathbb{E}_8$ and $\mathbb{E}_7$ respectively. For a lattice M and a rational t , we denote by $M(t)$ the lattice which is obtained by multiplication by t of the form of M .

Assume $\delta_F = 1$. Then $r \geq 16$. By Figure 6, we should check the genus invariants $(r, a, \delta_\varphi) = (18, 2, 0)$, $(17, 1, 1)$ and $(16, 2, 1)$.

If $(r, a, \delta_\varphi) = (18, 2, 0)$, then $(L_1)_{\varphi_1}$ is an even unimodular hyperbolic lattice of signature $(1, 1)$, and $h \in (L_1)_{\varphi_1}$ is a primitive element with $h^2 = -4$. Then $K = h^\perp$ in $(L_1)_{\varphi_1}$ is isomorphic to $\langle 4 \rangle$, and (3.4) is obvious.

If $(r, a, \delta_\varphi) = (17, 1, 1)$ then $(L_1)^{\varphi_1} \cong U \oplus E_8 \oplus E_7$, and K is orthogonal complement to $(L_1)^{\varphi_1} \oplus \mathbb{Z}h$ in L_1 . Then K has the genus of $\langle 4 \rangle \oplus \langle -2 \rangle$. Equivalently, $K(1/2)$ has the genus of $\langle 2 \rangle \oplus \langle -1 \rangle$. Thus, $K(1/2)$ can be obtained as odd orthogonal complement to an element $\delta \in M$ with $\delta^2 = -2$ where M is the odd unimodular lattice $\langle 1 \rangle \oplus 2\langle -1 \rangle$.

If $(r, a, \delta) = (16, 2, 1)$, then the same considerations show that $K(1/2)$ can be obtained as odd orthogonal complement to an element $\delta \in M$ with $\delta^2 = -2$ where $M = \langle 1 \rangle \oplus 3\langle -1 \rangle$.

Assume $\delta_F = 0$. Then $r \geq 17$, and the genus invariants of Figure 5 give $(r, a, \delta_\varphi) = (18, 2, 0)$, $(17, 3, 1)$.

If $(r, a, \delta_\varphi) = (18, 2, 0)$, then the lattice $(L_1)_{\varphi_1} \cong U(2)$, and K is the orthogonal complement to $h \in U(2)$ with $h^2 = -4$. Then $K \cong \langle 4 \rangle$, and (3.4) is obvious.

If $(r, a, \delta_\varphi) = (17, 3, 1)$, then $(L_1)_{\varphi_1} \cong \langle 2 \rangle \oplus 2\langle -2 \rangle$. It follows that $K(1/2)$ is odd orthogonal complement to an element $h \in \langle 1 \rangle \oplus 2\langle -1 \rangle$ with $h^2 = -2$.

Thus, for a lattice $M = \langle 1 \rangle \oplus t\langle -1 \rangle$ where $t = 2$ or $t = 3$ we have to check the following statement: The odd orthogonal complements K to elements $\delta \in M$ with $\delta^2 = -2$ are all isomorphic, and the canonical homomorphism $O(K) \rightarrow O(q_{K(2)})$ is epimorphic. Exactly this statement had been checked in the proof of Theorem 21 of [19].

It follows the statement. $\square$

Thus, we finally enumerated connected components of moduli we are looking for, and we obtained the main result.

Theorem 13. *There are exactly 108 connected components of moduli of real non-singular curves $A \in |-2K_{\mathbb{F}_2}|$ up to $\text{Aut}(\mathbb{F}_2/\mathbb{R})$, equivalently of related pairs of positive real non-singular curves $A \in |-2K_{\mathbb{F}_2}|$, equivalently of related pairs of connected components of moduli of real K3 surfaces with degenerate non-symplectic involution of type $(2, 2, 0)$.*

They are labelled by their invariants (3.3) satisfying relations 0.1–7, I.1–8, enumerated in Figures 5 and 6.

In [19] some other invariants of real K3 surfaces with non-symplectic involution of type $(2, 2, 0)$ were used. For further considerations and applications it is important to rewrite invariants (3.3) in terms of that invariants.

For an involution (L, τ, φ) from (3.1) and $S = [C, F] = L^\tau$ we define the following invariants. First, like δ_F , for any $x_- \in S$ we can define $\delta_{x_-} \in \{0, 1\}$. Here $\delta_{x_-} = 0$ if and only if $x_- \cdot L_\varphi \equiv 0 \pmod{2}$. For example, $\delta_C = 1$ because $C \cdot F_1 = 1$. Similarly $\delta_{C+F} = 1$ because $(C + F) \cdot F_1 = -1$. Thus, only δ_F can take two values: 0 or 1.

Now we define a subgroup $H \subset S/2S$ by the condition:

$$(3.5) \quad x_- \pmod{2S} \in H \text{ iff } \delta_{x_-} = 0.$$

Thus, for our (L, τ, φ) we have $H = \{0\}$ if $\delta_F = 1$, and $H = \mathbb{Z}/2\mathbb{Z}cl(F)$ if $\delta_F = 0$.

We say that $\delta_{\varphi S} = 0$ if there exists $v \in S \pmod{2S}$ such that $x \cdot \varphi(x) \equiv x \cdot v \pmod{2}$ for any $x \in L$. Otherwise, $\delta_{\varphi S} = 1$. If $\delta_{\varphi S} = 0$, then the element $v \pmod{2S}$ is defined uniquely, and it is called the *characteristic element of φ* . If $\delta_{\varphi S} = 1$, then the characteristic element v of φ is not defined. It is easy to see that if $\delta_{\varphi S} = 0$ and v is the characteristic element, then $\delta_v = 0$.

In our case, we then have

$$(\delta_{\varphi S} = 0, v = 0) \text{ iff } \delta_\varphi = 0;$$

and

$$(\delta_{\varphi S} = 0, v = cl(F)) \text{ iff } \delta_{\varphi F} = 0;$$

$$\delta_{\varphi S} = 1 \text{ iff } \delta_\varphi = \delta_{\varphi F} = 1.$$

Thus, all invariants (3.3) are equivalent to

$$(3.6) \quad (r, a, H, \delta_{\varphi S}, v)$$

where v is defined only for $\delta_{\varphi S} = 0$.

The invariants (3.6) are important because the same invariants are defined for the related involution $\tilde{\varphi} = \tau\varphi$ by replacing everywhere φ by $\tilde{\varphi}$. In [19] their relation had been found. Let us denote by a_H the rank of H over $\mathbb{Z}/2\mathbb{Z}$. On $S/2S$ the non-degenerate pairing b_S is defined by the condition $b_S(x + 2S, y + 2S) = x \cdot y/2 \pmod{2}$. It is skew-symmetric: $b_S(x + 2S, x + 2S) = 0 \pmod{2}$. We have (see (4.2) and (4.3) in [19]):

$$(3.7) \quad \begin{aligned} &\varphi(cl(F_1)) = -cl(F_1), \varphi(cl(F_2)) = -cl(F_2); \tilde{\varphi}(cl(F_1)) = -cl(F_2), \tilde{\varphi}(cl(F_2)) = -cl(F_1); \\ &r(\varphi) + r(\tilde{\varphi}) = 20; a_{H(\varphi)} + a_{H(\tilde{\varphi})} = 2; a(\varphi) - a_{H(\varphi)} = a(\tilde{\varphi}) - a_{H(\tilde{\varphi})}; \\ &H(\tilde{\varphi}) = H(\varphi)^\perp \text{ with respect to } b_S; \delta_{\tilde{\varphi}S} = \delta_{\varphi S}; v(\tilde{\varphi}) = v(\varphi). \end{aligned}$$

Thus, $H(\tilde{\varphi}) = [F] = (\mathbb{Z}/2\mathbb{Z})cl(F)$ if $\delta_F(\varphi) = 0$, and $H(\tilde{\varphi}) = [S] = S/2S$ if $\delta_F(\varphi) = 1$. Applying (3.7), from invariants of Figures 5, 6 we obtain all invariants of $\tilde{\varphi}$.

Below we consider geometric interpretation of the invariants (3.3) in terms of K3 and curves on $\mathbb{F}_2$.

3.3. Geometric interpretation of the invariants (3.3) in terms of K3. For the anti-holomorphic involutions φ and $\tilde{\varphi}$ we have the same geometric interpretations as in Sect. 2.3 for $\mathbb{F}_3$. One should only use the invariants $r(\varphi)$, $a(\varphi)$, δ_φ , $\delta_x(\varphi)$, $\delta_{x\varphi}$, $x \in S$, for φ , and $r(\tilde{\varphi})$, $a(\tilde{\varphi})$, $\delta_{\tilde{\varphi}}$, $\delta_x(\tilde{\varphi})$, $\delta_{x\tilde{\varphi}}$, for $\tilde{\varphi}$. For example, $\delta_{\tilde{\varphi}C} = 0$ if and only if $X_{\tilde{\varphi}}(\mathbb{R}) \sim C(\mathbb{C}) \pmod{2}$ in $H_2(X(\mathbb{C}); \mathbb{Z})$.

Using results above, like for $\mathbb{F}_2$ we then obtain

Theorem 14. *The connected component of moduli of a real K3 surface with degenerate non-symplectic involution (X, τ, φ) (or $(X, \tau, \tilde{\varphi})$) of type $(2, 2, 0)$, equivalently of a positive real non-singular curve $A \in |-2K_{\mathbb{F}_2}|$, is determined by the geometric invariants: topology of the real part $X_\varphi(\mathbb{R})$, if $F_\varphi(\mathbb{R}) \sim 0 \pmod{2}$ in $H_1(X_\varphi(\mathbb{R}); \mathbb{Z})$, if $X_\varphi(\mathbb{R}) \sim 0 \pmod{2}$ in $H_2(X(\mathbb{C}); \mathbb{Z})$, if $X_\varphi(\mathbb{R}) \sim F(\mathbb{C}) \pmod{2}$ in $H_2(X(\mathbb{C}); \mathbb{Z})$.*

Theorem 15. *The connected component of moduli of a real K3 surface with degenerate non-symplectic involution (X, τ, φ) (or $(X, \tau, \tilde{\varphi})$) of type $(2, 2, 0)$, equivalently of a positive real non-singular curve $A \in |-2K_{\mathbb{F}_2}|$, is determined by the geometric invariants: topology of the real part $X_{\tilde{\varphi}}(\mathbb{R})$, if $C_{\tilde{\varphi}}(\mathbb{R}) \sim 0 \pmod{2}$ in $H_1(X_{\tilde{\varphi}}(\mathbb{R}); \mathbb{Z})$, if $X_{\tilde{\varphi}}(\mathbb{R}) \sim 0 \pmod{2}$ in $H_2(X(\mathbb{C}); \mathbb{Z})$, if $X_{\tilde{\varphi}}(\mathbb{R}) \sim F(\mathbb{C}) \pmod{2}$ in $H_2(X(\mathbb{C}); \mathbb{Z})$.*

3.4. Geometric interpretation of the invariants (3.3) in terms of $A \in |-2K_{\mathbb{F}_2}|$. By Theorem 5, the real curve $A = X^\tau$ is dividing, i. e. $cl(A(\mathbb{R})) = 0$ in $H_1(A(\mathbb{C}); \mathbb{Z}/2\mathbb{Z})$, if and only if $\delta_{\varphi S} = 0$ (equivalently, $\delta_{\tilde{\varphi} S} = 0$).

By results above, $X_\varphi(\mathbb{R})$ and $X_{\tilde{\varphi}}(\mathbb{R})$ are both empty and then $\mathbb{F}_2(\mathbb{R})$ is empty if and only if the invariants (3.3) are: $\delta_F = 0$ and $(r, a, \delta_\varphi) = (10, 10, 0)$. This gives only one connected component of moduli of non-singular curves $A \in |-2K_{\mathbb{F}_2}|$, and two related connected components of moduli of positive curves.

Further we assume that $\mathbb{F}_2(\mathbb{R}) \neq \emptyset$, equivalently, the invariants (3.3) are different from $(\delta_F, r, a, \delta_\varphi) = (0, 10, 10, 0)$. See Figures 5 and 6. Then $s(\mathbb{R}) \neq \emptyset$, and the rational pencil of $\mathbb{F}_2$ gives a pencil over the circle $s(\mathbb{R})$ by circles $c(\mathbb{R})$. Further we always take the rational fibre to be real, i. e. over $s(\mathbb{R})$. Since $s^2 = -2$ is even, it follows that $\mathbb{F}_2(\mathbb{R})$ is a torus where $s(\mathbb{R})$ and $c(\mathbb{R})$ are generators of the torus (i. e. of $H_1(\mathbb{F}_2(\mathbb{R}); \mathbb{Z})$).

A component of $A(\mathbb{R})$ is called *oval* if it bounds a disk in $\mathbb{F}_2(\mathbb{R})$. Otherwise, it is called non-oval.

We always consider the positive curve $A^+ = \pi(X_\varphi(\mathbb{R}))$, i. e. when $s(\mathbb{R}) \subset A^+$. Since A does not intersect s and $c \cdot A = 4$, by elementary 2-dimensional topology considerations, we have the following and only the following pictures where we use the geometric interpretation of invariants (3.3) above in terms of the double covering K3. Moreover, to draw correctly $s(\mathbb{R})$ in Figures 10 and 11, we use ellipsoid deformation from Sect. 3.7 below.

Case 1. $A(\mathbb{R})$ has no non-ovals in $\mathbb{F}_2(\mathbb{R}) \neq \emptyset$. Then $\delta_F = 1$ for the invariants (3.3), and the positive curve $A^+ = \pi(X_\varphi(\mathbb{R}))$ is shown (as shadow) in Figures 7, 8 and 9. All invariants (3.3) with $\delta_F = 1$ are covered by this case.

Case 2. $A(\mathbb{R})$ has non-ovals. Then $\delta_F = 0$ for the invariants (3.3), and the positive curve $A^+ = \pi(X_\varphi(\mathbb{R}))$ is shown in Figures 10, 11 and 12. This case covers all invariants (3.3) with $\delta_F = 0$ except $(r, a, \delta_\varphi) = (10, 10, 0)$ when $\mathbb{F}_2(\mathbb{R}) = \emptyset$. See Figure 5 for the full list of these invariants. If $\delta_F = 0$ and the invariants $(r, a, \delta_\varphi) \neq (10, 10, 0)$ nor $(10, 8, 0)$, then $A(\mathbb{R})$ has exactly two non-ovals, and A^+ is shown in Figures 10, 11. If $(r, a, \delta_\varphi) = (10, 8, 0)$, then $A(\mathbb{R})$ consists of four non-ovals, and A^+ is shown in Figure 12.

Thus, we obtain

Theorem 16. *The connected component of moduli of a positive curve A^+ of a real non-singular curve $A \in |-2K_{\mathbb{F}_2}|$ is determined by the following invariants: the isotopy type of A^+ in $\mathbb{F}_2(\mathbb{R})$, by dividedness by $A(\mathbb{R})$ of $A(\mathbb{C})$, if $s(\mathbb{R}) \subset A^+$. If $\mathbb{F}_2(\mathbb{R}) = \emptyset$, there are exactly two related connected components of moduli.*




FIGURE 7. $\mathbb{F}_2$: The positive curve $A^+ = \pi(X_\varphi(\mathbb{R}))$ for $\delta_F = 1$ and $(r, a, \delta_\varphi) \neq (10, 8, 0), (10, 6, 0)$ (see Figure 6), here $g = (22 - r - a)/2$, $k = (r - a)/2$.




FIGURE 8. $\mathbb{F}_2$: The positive curve $A^+ = \pi(X_\varphi(\mathbb{R}))$ for $\delta_F = 1$ and $(r, a, \delta_\varphi) = (10, 6, 0)$ (see Figure 6).

All possibilities are presented in Figures 5 —12 and 3.7.




FIGURE 9. $\mathbb{F}_2$: The positive curve $A^+ = \pi(X_\varphi(\mathbb{R}))$ for $\delta_F = 1$ and $(r, a, \delta_\varphi) = (10, 8, 0)$ (see Figure 6). Then $X_\varphi(\mathbb{R}) = (T_1)^2$ and $A(\mathbb{R}) = \emptyset$.




FIGURE 10. $\mathbb{F}_2$: The positive curve $A^+ = \pi(X_\varphi(\mathbb{R}))$ for $\delta_F = 0$ and $(r, a, \delta_\varphi) \neq (10, 10, 0)$, $(10, 8, 0)$, and $(r, a, \delta_{\varphi_F}) \neq (8, 8, 0)$ (see Figure 5), here $g = (22 - r - a)/2$, $k = (r - a)/2$.

3.5. Application: Connected components of moduli of real non-singular degree 8 curves on a quadratic cone in $\mathbb{P}^3$. We have $h_1 = -K_{\mathbb{F}_2}/2 = 2c + s$ has $h_1^2 = 2$. The linear system $|h_1|$ gives the embedding of $\mathbb{F}_2$ as a quadratic cone in $\mathbb{P}^3$ with the exceptional section s contracted to the vertex of the cone. The linear system $|-2K_{\mathbb{F}_2}|$ consists then of




FIGURE 11. $\mathbb{F}_2$: The positive curve $A^+ = \pi(X_\varphi(\mathbb{R}))$ for $\delta_F = 0$ and $(r, a, \delta_{\varphi F}) = (8, 8, 0)$ (see Figure 5).




FIGURE 12. $\mathbb{F}_2$: The positive curve $A^+ = \pi(X_\varphi(\mathbb{R}))$ for $\delta_F = 0$ and $(r, a, \delta_\varphi) = (10, 8, 0)$ (see Figure 5).

sections of the cone by the degree 4 forms, i. e. of the degree 8 curves, which do not pass through the vertex of the cone.

Let $(\xi_0 : \xi_1 : \xi_2 : \xi_3)$ be usual real homogeneous coordinates of $\mathbb{P}^3$. The cone is given by the equation $Q_2 = \xi_1^2 + \xi_2^2 - \xi_3^2 = 0$ if $\mathbb{F}_2(\mathbb{R}) \neq \emptyset$, and by the equation $\xi_1^2 + \xi_2^2 + \xi_3^2 = 0$ if $\mathbb{F}_2(\mathbb{R}) = \emptyset$. The real curve A is given by the equation $F_4 = 0$ on the cone $Q_2 = 0$ which must

be non-singular as a complex algebraic curve. Here $F_4(\xi_0, \xi_1, \xi_2, \xi_3)$ is a homogeneous degree four form with real coefficients which is defined up to $\lambda(F_4 + Q_2 F_2)$ where $\lambda \in \mathbb{R}^*$ and F_2 a degree two form $F_2(\xi_0, \xi_1, \xi_2, \xi_3)$ with real coefficients. The positive curve A^+ is given by the inequality $F_4 \geq 0$ (or $-F_4 \geq 0$) on the cone such that A^+ contains the vertex $(1 : 0 : 0 : 0)$ of the cone.

Thus, we obtain classification of connected components of moduli of real non-singular degree 8 curves on the real cone in $\mathbb{P}^3$ up to the automorphism group of the cone which is the group $PO(Q_2)$. It consists of projective real transformations which multiply the form Q_2 by a constant.

We leave further geometric interpretation of our results above in terms of degree 8 curves on a quadratic cone to the interested reader.

It is important and interesting to consider deformations of non-singular degree 8 curves on a quadratic cone to non-singular degree 8 curves on a non-degenerate quadric: One replaces the degenerate quadratic form Q_2 defining the cone $Q_2 = 0$ (which is $\mathbb{F}_2$ with contracted s) by a closed non-degenerate quadratic form Q'_2 , and the degree four form F_4 defining a non-singular curve A on the cone ($Q_2 = F_4 = 0$) by a closed degree four form F'_4 . Then equations $Q'_2 = F'_4 = 0$ define a non-singular bidegree $(4, 4)$ curve A' on the non-degenerate quadric $Q' = 0$ which is $\mathbb{F}_0$.

There are two very different types of these deformations. First type when the quadric $Q'_2 = 0$ is different from ellipsoid (it is a hyperboloid or empty), e. g. the form $Q_2 = \xi_1^2 + \xi_2^2 - \xi_3^2$ is replaced by the form $-\varepsilon \xi_0^2 + \xi_1^2 + \xi_2^2 - \xi_3^2$ where $\varepsilon > 0$ is small. Second type when the quadric $Q'_2 = 0$ is ellipsoid, e. g. the form $Q_2 = \xi_1^2 + \xi_2^2 - \xi_3^2$ is replaced by the form $Q'_2 = \varepsilon \xi_0^2 + \xi_1^2 + \xi_2^2 - \xi_3^2$ where $\varepsilon > 0$ is small. Second type deformations, when the quadric $Q'_2 = 0$ is ellipsoid, is called *ellipsoid deformation*. All other are called *hyperboloid deformations*.

Further we study these deformations in details.

3.6. Application: Hyperboloid deformations of $A \in |-2K_{\mathbb{F}_2}|$ to $A \in |-2K_{\mathbb{F}_0}|$. Let us consider a real K3 surface with degenerate non-symplectic involution (X, τ, φ) (or $(X, \tau, \tilde{\varphi})$) of type $(2, 2, 0)$. It defines then a connected component of moduli of real K3 surfaces with non-symplectic involution of type $(2, 2, 0)$ where we consider non-degenerate and degenerate involutions all together. By [19], the connected component of moduli is also defined by the isomorphism class of the action (L, τ, φ) with the condition $\varphi = -1$ on $S = L^\tau = [C, F = F_1 + F_2]$ for the hyperboloid case (for ellipsoid case it is different, see below). We just forget about F_1 and F_2 , and are interested only on their sum $cl(F) = cl(F_1) + cl(F_2)$. Moreover, connected components of moduli of all involutions and non-degenerate ones coincide.

In [19] other generators for $S = L^\tau$ (related with $\mathbb{F}_0$) were used. They are $e_1 = cl(C)$, $e_2 = cl(C) + cl(F)$ (i. e. the effective isotropic generators of S corresponding to the structure $\mathbb{F}_0 = \mathbb{P}^1 \times \mathbb{P}^1$) and $h = e_1 + e_2 = 2cl(C) + cl(F)$ (which is pre-image of hyperplane of $\mathbb{P}^3$). Thus $cl(F) \equiv e_1 + e_2 = h \pmod{2}$. Invariants of connected components of moduli for hyperboloid case are the same as we used in Sect. 3.2. See [19]. Thus, hyperbolic deformations of non-singular curves $A \in |-2K_{\mathbb{F}_2}|$ to non-singular curves $A \in |-2K_{\mathbb{F}_0}|$ can be described

symbolically as follows:

$$(3.8) \quad \begin{aligned} (\mathbb{F}_2; r(\varphi), a(\varphi), H(\varphi), \delta_{\varphi S}, v(\varphi)) &\implies (\mathbb{H} : r = r(\varphi), a = a(\varphi), H = H(\varphi), \delta_{\varphi S}, v = v(\varphi)); \\ (\mathbb{F}_2; r(\tilde{\varphi}), a(\tilde{\varphi}), H(\tilde{\varphi}), \delta_{\tilde{\varphi} S}, v(\tilde{\varphi})) &\implies (\mathbb{H} : r = r(\tilde{\varphi}), a = a(\tilde{\varphi}), H = H(\tilde{\varphi}), \delta_{\tilde{\varphi} S}, v = v(\tilde{\varphi})). \end{aligned}$$

where to the left we have invariants of a connected component of moduli of positive real non-singular curves on $\mathbb{F}_2$, and to the right we have invariants of a connected component of moduli of positive real non-singular curves on hyperboloid (see [19]) obtained under hyperboloid deformation.

Using (3.8), we can answer the following interesting question. When there exist such deformation, and how many of them do exist? Comparing invariants of connected components for hyperboloid from [19] and for $\mathbb{F}_2$ from Sect. 3.2, we obtain the result.

For $\mathbb{F}_2$, the groups $H(\varphi)$ and $H(\tilde{\varphi})$ cannot be equal to $\mathbb{Z}/2\mathbb{Z}e_1$ or $\mathbb{Z}/2\mathbb{Z}e_2$. The group $H(\varphi)$ cannot be $S/2S$. The group $H(\tilde{\varphi})$ cannot be zero. The invariants $(r = 12, a = 8, H = [e_1 + e_2], \delta_{\varphi S} = 0, v = e_1 + e_2)$ of hyperboloid cannot be obtained from φ on $\mathbb{F}_2$ (see Figure 5) but they can be obtained from $\tilde{\varphi}$ on $\mathbb{F}_2$. Similarly, the invariants $(r = 8, a = 8, H = [e_1 + e_2], \delta_{\varphi S} = 0, v = e_1 + e_2)$ of hyperboloid can be obtained from φ on $\mathbb{F}_2$, but they cannot be obtained from $\tilde{\varphi}$ on $\mathbb{F}_2$. Using Sect. 3.4 and geometric interpretation of the invariants for hyperboloid from [19], we can formulate the result in geometric terms. Fortunately, here we don't have too many possibilities.

Theorem 17. *If $\mathbb{F}_0/\mathbb{R}$ is different from ellipsoid, then a positive real non-singular curve $A \in |-2K_{\mathbb{F}_0}|$ can be obtained by deformation of a positive real non-singular curve $A_0 \in |-2K_{\mathbb{F}_2}|$ if and only if $\mathbb{F}_0/\mathbb{R}$ is hyperboloid and $A(\mathbb{R})$ does not have connected components homological to any line (l_1 or l_2 from different generators) of hyperboloid. See Figures 10–13 in [19].*

This deformation is unique if either $A(\mathbb{R})$ consists of only ovals (including empty $A(\mathbb{R})$) on hyperboloid (see Figures 10, 11 in [19]), or $A(\mathbb{R})$ consists of two non-ovals homological to $l_1 + l_2$ and two ovals between these non-ovals (then A^+ is connected) and A is dividing. (See Figure 12 in [19] for $g = 3, k = 0$ or $g = 1, k = 2$.) In the last case, the connected component of the positive curve of A_0 containing two ovals, must also contain $s(\mathbb{R})$.

Otherwise, there are exactly two isomorphism classes of such deformations: for one of them $s(\mathbb{R})$ is contained in the positive curve of A_0 , and for another one $s(\mathbb{R})$ is not contained in the positive curve of A_0 .

3.7. Application: Ellipsoid deformations of $A \in |-2K_{\mathbb{F}_2}|$ to $A \in |-2K_{\mathbb{F}_0}|$. We describe the ellipsoid deformation of a positive real non-singular curve $A \in |-2K_{\mathbb{F}_2}|$ in terms of periods of K3 surfaces using Global Torelli Theorem for K3 surfaces [20].

For an $A \in |-2K_{\mathbb{F}_2}|$, let (X, τ, φ) be the corresponding real K3 surface with degenerate non-symplectic involution of type $(2, 2, 0)$, and (L, τ, φ) its action.

An element δ of a lattice M with $\delta^2 = -2$ defines the reflection s_δ of the lattice M by the formula $x \rightarrow x + (x \cdot \delta)\delta$, $x \in M$. We have $s_\delta(\delta) = -\delta$ and δ is identical in $\delta^\perp$.

Since $F_1^2 = F_2^2 = -2$, they define reflections $s_{cl(F_1)}$ and $s_{cl(F_2)}$ of L . Since $F_1 \cdot F_2 = 0$, the reflections $s_{cl(F_1)}$ and $s_{cl(F_2)}$ commute, and we obtain the involution

$$s_{[F_1, F_2]} = s_{cl(F_1)} s_{cl(F_2)}$$

of L with the eigenvalue -1 part $[F_1, F_2]$.

The involution $s_{[F_1, F_2]}$ obviously commutes with τ and φ . Let us replace the involutions φ and $\tilde{\varphi}$ on L by the involutions (we apply twice the Lefschetz formula for the vanishing cycle)

$$(3.9) \quad \psi = s_{[F_1, F_2]} \varphi, \quad \tilde{\psi} = s_{[F_1, F_2]} \tilde{\varphi} = \tau \psi.$$

They coincide with $\varphi, \tilde{\varphi}$ in $[F_1, F_2]^\perp$ respectively. This geometrically means that ψ, φ and $\tilde{\psi}, \tilde{\varphi}$ are the same outside the exceptional curves F_1 and F_2 which must take place for ellipsoid deformation.

The involution ψ coincides with φ in $[C, F_1, F_2]^\perp$, and $\psi(cl(F_1)) = cl(F_1), \psi(cl(F_2)) = cl(F_2)$. As above, $h = 2cl(C) + cl(F)$ generates the orthogonal complement to F in $S = [C, F]$, and $\psi(h) = -h, \psi(cl(F)) = cl(F)$. Thus, ψ acts in $S = L^\tau$ as ellipsoid involution (see [19]), and (L, τ, ψ) defines the connected component of moduli of curves $A \in |-2K_{\mathbb{F}_0}|$ where $\mathbb{F}_0$ is ellipsoid, see [19].

For degenerate (X, τ, φ) its periods (e. g. see [15] and Sect. 2.1 in [19]) are equivalent to a pair of rays $(\mathbb{R}^+ \omega_+, \mathbb{R}^+ \omega_-)$ where $\omega_+ \in L^\varphi \otimes \mathbb{R}$ and $\omega_+^2 > 0$, and $\omega_- \in L_\varphi \otimes \mathbb{R}$ where $\omega_-^2 > 0$ and $\omega_- \perp [C, F_1, F_2]$. Since $[C, F_1, F_2] \subset L_\varphi$, it follows that $\omega_- \in ([C, F_1, F_2]^\perp_{L_\varphi} \otimes \mathbb{R})$.

For ψ we have

$$L^\psi \otimes \mathbb{R} = L^\varphi \otimes \mathbb{R} \oplus \mathbb{R}(cl(F_1) - cl(F_2)) \oplus \mathbb{R}cl(F)$$

and

$$L_\psi \otimes \mathbb{R} = [C, F_1, F_2]^\perp_{L_\varphi} \otimes \mathbb{R} \oplus \mathbb{R}h.$$

Their orthogonal complements to $S = [C, F]$ consist of $L^\varphi \otimes \mathbb{R} \oplus \mathbb{R}(cl(F_1) - cl(F_2))$ and $[C, F_1, F_2]^\perp_{L_\varphi} \otimes \mathbb{R}$ respectively. For the ellipsoid deformation with the action (L, τ, ψ) , we must take for periods of real K3

$$\tilde{\omega}_+ \in (L^\varphi \otimes \mathbb{R} \oplus \mathbb{R}(cl(F_1) - cl(F_2))) - L^\varphi \otimes \mathbb{R}$$

instead of $\omega_+ \in L^\varphi \otimes \mathbb{R}$ (here $L^\varphi \otimes \mathbb{R} \subset (L^\varphi \otimes \mathbb{R} \oplus \mathbb{R}(cl(F_1) - cl(F_2)))$ has codimension one), and

$$\tilde{\omega}_- \in [C, F_1, F_2]^\perp_{L_\varphi} \otimes \mathbb{R}$$

which is the same space as for $\omega_- \in [C, F_1, F_2]^\perp_{L_\varphi} \otimes \mathbb{R}$. They can be taken closed to ω_+ and ω_- and correspond to periods of a real K3 surface with non-symplectic involution of ellipsoid type by Global Torelli Theorem for K3 surfaces [20].

Thus, for ellipsoid deformation, we must just replace (L, τ, φ) by (L, τ, ψ) where ψ is given by (3.9). For $(L, \tau, \tilde{\varphi})$ we must replace ψ by the related involution $\tilde{\psi} = \tau \psi$ which is given by the same formula (3.9).

In [19], for an ellipsoid involution (L, τ, ψ) the same invariants ($r = r(\psi), a = a(\psi), \delta = \delta_{\psi S}$) were used (the invariant $v = v(\psi)$ follows from these invariants). They define the connected component of moduli of a real K3 involution of ellipsoid type and corresponding positive real non-singular bidegree $(4, 4)$ curve on ellipsoid. It is easy to calculate these invariants. The main formula is

$$(3.10) \quad r(\tilde{\psi}) = r(\tilde{\varphi}), \quad a(\tilde{\psi}) = a(\tilde{\varphi}), \quad \delta_{S\tilde{\psi}} = \delta_{S\tilde{\varphi}}$$

(it is geometrically obvious). The remaining relations follow from

$$(3.11) \quad r(\psi) + r(\tilde{\psi}) = 22, \quad a(\psi) = a(\tilde{\psi}), \quad \delta_{\psi S} = \delta_{\tilde{\psi} S}$$




FIGURE 13. $\mathbb{E}$: All possible (r, a, δ) for ellipsoid.

(see formulae (5.2) and (5.3) in [19]) and (3.7). We obtain

$$(3.12) \quad r(\psi) = r(\varphi) + 2, \quad a(\psi) = a(\varphi) + 2\delta_F, \delta_{\psi S} = \delta_{\varphi S}, \quad v(\psi) = v(\varphi) + cl(F) \pmod{2}.$$

Here (3.12) follows from Theorem 11 from [19] applied to $\tau = -s_{[F_1, F_2]}$ and φ , which gives the formal proof of formulae (3.12) and (3.10).

Thus, the ellipsoid deformation can be represented symbolically as follows:

$$(3.13) \quad \begin{aligned} (\mathbb{F}_2 : r(\varphi), a(\varphi), \delta_F(\varphi) = 0, \delta_{\varphi S}, v(\varphi)) &\implies (\mathbb{E} : r = r(\varphi) + 2, a = a(\varphi), \delta = \delta_{\varphi S}); \\ (\mathbb{F}_2 : r(\varphi), a(\varphi), \delta_F(\varphi) = 1, \delta_{\varphi S}, v(\varphi)) &\implies (\mathbb{E} : r = r(\varphi) + 2, a = a(\varphi) + 2, \delta = \delta_{\varphi S}); \\ (\mathbb{F}_2 : r(\tilde{\varphi}), a(\tilde{\varphi}), H(\tilde{\varphi}) = [F], \delta_{\tilde{\varphi} S}, v(\tilde{\varphi})) &\implies (\mathbb{E} : r = r(\tilde{\varphi}), a = a(\tilde{\varphi}), \delta = \delta_{\tilde{\varphi} S}); \\ (\mathbb{F}_2 : r(\tilde{\varphi}), a(\tilde{\varphi}), H(\tilde{\varphi}) = [S], \delta_{\tilde{\varphi} S}, v(\tilde{\varphi})) &\implies (\mathbb{E} : r = r(\tilde{\varphi}), a = a(\tilde{\varphi}), \delta = \delta_{\tilde{\varphi} S}). \end{aligned}$$

There are four types of deformations.

All invariants (r, a, δ) for ellipsoid are presented in Figure 13 (it is Figure 16 from [19]).

From (3.13) and Figures 5, 6, 13 and (3.7) we see all deformations of curves from $\mathbb{F}_2$ to ellipsoid. It is hard to formulate similar to Theorem 17 result because there are too many cases. We restrict by the following.

Theorem 18. *Any positive real non-singular bidegree $(4, 4)$ curve on ellipsoid can be obtained by ellipsoid deformation of a positive real non-singular curve in $|-2K_{\mathbb{F}_2}|$. There are four isomorphism classes of such deformations at most. All of them are presented in (3.13), (3.7) and Figures 13, 5, 6.*

3.8. Application: Classification of connected components of moduli of real hyper-elliptically polarized K3 surfaces. For a positive real non-singular curve $A \in |-2K_Y|$ where $Y = \mathbb{F}_2$, or for the corresponding real K3 surface with degenerate non-symplectic involution (X, τ, φ) of type $(2, 2, 0)$, let us consider $P = n_1 C + F$ for $n_1 \geq 3$. Here P is primitive and has the degree $n = P^2 = 4n_1 - 4$ where $n \geq 8$ and $n \equiv 0 \pmod{4}$. Thus, $n_1 = (n/4) + 1$ and $P = ((n/4) + 1)C + F$. We have $C \cdot P = 2$ for the elliptic pencil $|C|$ on X . Then the complete linear system $|P|$ is hyper-elliptic and gives a double covering $|P| : X \rightarrow Y = \mathbb{F}_2 \subset \mathbb{P}^N$ where $N = P^2/2 + 1 = n/2 + 1$. Here the linear system $|P|$ is pre-image $\pi^*[n_1 c + s]$ of the standard complete linear system $|n_1 c + s|$ from Y . On the other hand, P is ample since $|P|$ gives a double covering ramified in a non-singular curve.

Thus, (X, P) is a real hyper-elliptically polarized K3 surface. See [22], [19] and [18] for more details.

Thus, our classification of the connected components of moduli also enumerates connected components of moduli of real hyper-elliptically polarized K3 surfaces (X, P) of that type. We shall call them as *hyper-elliptically polarized K3 surfaces of $\mathbb{F}_2$ type*.

All other types had been considered and classified in [19]. See [19] and [18] for details. For other types, one has to take $Y = \mathbb{P}^2$ or $\mathbb{F}_m$ with $m = 0, 1, 4$ instead of $\mathbb{F}_2$ and also pick up some standard linear system from Y . All these possibilities were described in Sect. 8 of [19] (see also [18]). Thus, here we finished classification of connected components of moduli of real hyper-elliptically polarized K3 surfaces describing the last $\mathbb{F}_2$ type which had not been considered. We remark that for a general hyper-elliptically polarized K3 surface of $\mathbb{F}_2$ type the Picard number is 3, and then the dimension of moduli is 17. For all other types of hyper-elliptically polarized K3 surfaces the Picard number is 2 and the dimension of moduli is 18.

Thus, we obtain the following result.

Theorem 19. *Real hyper-elliptically polarized K3 surfaces (X, P) of type $\mathbb{F}_2$ (i. e. when the image $|P| : X \rightarrow Y$ is $Y = \mathbb{F}_2$) exist only for primitive P and $n = P^2 \geq 8$ where $n \equiv 0 \pmod{4}$. For each such n there are exactly 216 connected components of their moduli (of the dimension 17) which are enumerated by natural invariants in Theorem 16.*

Similar classification for all other types had been obtained in [19] (see also [18]). Thus, Theorem 19 completes this classification.

3.9. Application: Real polarized K3 surfaces as deformations of real hyper-elliptically polarized K3 surfaces of type $\mathbb{F}_2$. Let (X, P) be a real hyper-elliptically polarized K3 surface of type $\mathbb{F}_2$. Its small deformation gives then a polarized K3 surface $(\tilde{X}, \tilde{P})$ where $\tilde{P}$ is very ample and defines a connected component of moduli of real polarized K3 surfaces which are also deformations of (X, P) . Thus, we ask: What real polarized K3 surfaces are deformations of real hyper-elliptically polarized K3 surfaces of type $\mathbb{F}_2$? See [19] and [18] for further details about this problem where the same problem had been solved for all other types of real hyper-elliptically polarized K3 surfaces. Thus here we finalize these results.

By [15], the connected component of moduli of a real polarized K3 surface is determined by isomorphism class of the integral polarized involution (L, φ, P) corresponding to (X, τ, φ) with $P = (n/4 + 1)cl(C) + cl(F) \in S = L^\tau$ (or the same for $(X, \tau, \tilde{\varphi})$). Here $n \equiv 0 \pmod{4}$ and $n \geq 4$ (we can consider $n = 4$ as well, see below). The main invariants (the genus invariants) of the integral polarized involution (L, φ, P) are the same invariants as we used before:

$$(3.14) \quad (n; r = r(\varphi), a = a(\varphi), \delta_P; \delta_\varphi, \delta_{\varphi P})$$

Almost in all cases these invariants completely determine the connected component of moduli of real polarized K3 surfaces (see [15] and [18]). Thus, in our case there are four types of such deformations which can be symbolically expressed as follows where $n \geq 4$ and $n \equiv 0$

mod 4:

(3.15)

$$\begin{aligned} D_n : (\mathbb{F}_2; r(\varphi), a(\varphi), \delta_F(\varphi) = 1, \delta_\varphi, \delta_{\varphi F} = 1) &\implies (n; r = r(\varphi), a = a(\varphi), \delta_P = 1, \delta_\varphi, \delta_{\varphi P} = 1); \\ D_n : (\mathbb{F}_2; r(\varphi), a(\varphi), \delta_F(\varphi) = 0, \delta_\varphi, \delta_{\varphi F}) &\implies (n; r = r(\varphi), a = a(\varphi), \delta_P, \delta_\varphi, \delta_{\varphi P}) \text{ where} \end{aligned}$$

$$\delta_P = \begin{cases} 0 & \text{if } n \equiv 4 \pmod{8} \\ 1 & \text{otherwise} \end{cases}, \quad \delta_{\varphi P} = \begin{cases} 0 & \text{if } (n \pmod{8}, \delta_{\varphi F}) = (4 \pmod{8}, 0) \\ 1 & \text{otherwise} \end{cases};$$

$$\begin{aligned} D_n : (\mathbb{F}_2; r(\tilde{\varphi}), a(\tilde{\varphi}), H(\tilde{\varphi}) = [S], \delta_{\tilde{\varphi}}) &\implies (n; r = r(\tilde{\varphi}), a = a(\tilde{\varphi}), \delta_P = 0, \delta_{\tilde{\varphi}}, \delta_{\tilde{\varphi}P} = 1); \\ D_n : (\mathbb{F}_2; r(\tilde{\varphi}), a(\tilde{\varphi}), H(\tilde{\varphi}) = [F], \delta_{\tilde{\varphi}}, \delta_{\tilde{\varphi}F}) &\implies (n; r = r(\tilde{\varphi}), a = a(\tilde{\varphi}), \delta_P, \delta_{\tilde{\varphi}}, \delta_{\tilde{\varphi}P}) \text{ where} \end{aligned}$$

$$\delta_P = \begin{cases} 0 & \text{if } n \equiv 4 \pmod{8} \\ 1 & \text{otherwise} \end{cases}, \quad \delta_{\tilde{\varphi}P} = \begin{cases} 0 & \text{if } (n \pmod{8}, \delta_{\tilde{\varphi}F}) = (4 \pmod{8}, 0) \\ 1 & \text{otherwise} \end{cases}.$$

By [15] and [18], the invariants (3.14) determine the connected component of moduli of real polarized K3 surfaces if $r(\varphi) \leq 18$. Thus (3.15) completely determine deformations for $r(\varphi) \leq 18$ and $r(\tilde{\varphi}) \leq 18$.

By Figures 5 and 6, there are no cases with $r(\varphi) > 18$. By (3.7), $r(\tilde{\varphi}) > 18$ only for one connected component ($r(\tilde{\varphi}) = 19, a(\tilde{\varphi}) = 3, H(\tilde{\varphi}) = [S], \delta_{\tilde{\varphi}} = 1, \delta_{\tilde{\varphi}P} = 1$). Then $X_{\tilde{\varphi}}(\mathbb{R}) = (T_0)^9$. Thus, the deformation D_n can give only one connected component of moduli of real polarized K3 surfaces (for each $n \geq 4, n \equiv 0 \pmod{4}$). By Theorem 16 in [18], the connected component of moduli of real polarized K3 surfaces is *the standard connected component*.

The list of all possible invariants (3.14) of real polarized K3 surfaces is known, see [15] (and also [18]). Thus, using (3.15) and Figures 5, 6, and (3.7), we can find all real polarized K3 surfaces which are deformations of real hyper-elliptically polarized K3 surfaces of type $\mathbb{F}_2$. They must appear to the right of the deformations (3.15). We can also find out how many these deformations do exist. Three, at most. We leave further details to the interested reader.

We obtain the following.

Theorem 20. *All real polarized K3 surfaces which are deformations of real hyper-elliptically polarized K3 surfaces of type $\mathbb{F}_2$ and all isomorphism classes of these deformations are completely determined by the symbolical deformations (3.15), and Figures 5, 6 and (3.7) where $n \equiv 0 \pmod{4}$ and $n \geq 8$. For a fixed real polarized K3 surface there are three isomorphism classes of these deformations at most.*

For real hyper-elliptically polarized K3 surfaces of all other types, similar results were obtained in [19] and [18]. Thus, Theorem 20 finalizes these results by considering the last possible type.

Formally, Theorem 20 does not include $n = 4$ because $P = C + F$ is not ample for $n = 4$. We have $P \cdot F = 0$, and the linear system $|P|$ contracts $F_1 + F_2$.

But, considering hyperboloid or ellipsoid deformations $\mathbb{F}_2 \implies \mathbb{H}$ (see (3.8)), $\mathbb{F}_2 \implies \mathbb{E}$ (see (3.13)), and then taking $P = 2cl(C) + cl(F) = e_1 + e_2$, we obtain real hyper-elliptically polarized K3 surfaces of type $\mathbb{F}_0$ (over $\mathbb{R}$ they are of hyperboloid or ellipsoid type). Considering their deformations we obtain real polarized K3 surfaces with polarization of degree four (a non-singular quartic in $\mathbb{P}^3$, in general). These deformations were studied in [19]. Taking composition of these two deformations, we can consider $n = 4$ for $\mathbb{F}_2$ as well.

The hyperboloid deformation D_4 gives the same as D_4 in (3.15), and we obtain:

$$(3.16) \quad \begin{aligned} D_4 : (\mathbb{F}_2; r(\varphi), a(\varphi), \delta_F(\varphi) = 1, \delta_\varphi, \delta_{\varphi F} = 1) &\xrightarrow{\mathbb{H}} (4; r = r(\varphi), a = a(\varphi), \delta_P = 1, \delta_\varphi, \delta_{\varphi P} = 1); \\ D_4 : (\mathbb{F}_2; r(\varphi), a(\varphi), \delta_F(\varphi) = 0, \delta_\varphi, \delta_{\varphi F}) &\xrightarrow{\mathbb{H}} (4; r = r(\varphi), a = a(\varphi), \delta_P = 0, \delta_\varphi, \delta_{\varphi P} = \delta_{\varphi F}); \\ D_4 : (\mathbb{F}_2; r(\tilde{\varphi}), a(\tilde{\varphi}), H(\tilde{\varphi}) = [S], \delta_{\tilde{\varphi}}) &\xrightarrow{\mathbb{H}} (4; r = r(\tilde{\varphi}), a = a(\tilde{\varphi}), \delta_P = 0, \delta_{\tilde{\varphi}}, \delta_{\tilde{\varphi} P} = 1); \\ D_4 : (\mathbb{F}_2; r(\tilde{\varphi}), a(\tilde{\varphi}), H(\tilde{\varphi}) = [F], \delta_{\tilde{\varphi}}, \delta_{\tilde{\varphi} F}) &\xrightarrow{\mathbb{H}} (4; r = r(\tilde{\varphi}), a = a(\tilde{\varphi}), \delta_P = 0, \delta_{\tilde{\varphi}}, \delta_{\tilde{\varphi} P} = \delta_{\tilde{\varphi} F}). \end{aligned}$$

Obviously, all deformations D_n in (3.15) are compositions of deformations (3.8) of $\mathbb{F}_2$ to hyperboloid, and the deformation D_n of type $(\mathbb{H})_1$ (when $P = (n/4 + 1)C + F = (n/4)e_1 + e_2$ for hyperboloid) considered in [19]. (It had been used in [18].)

For ellipsoid deformation we obtain the following where we denote by (L, ψ, P) the integral polarized involution of a K3 surface which is obtained under deformation:

$$(3.17) \quad \begin{aligned} D_4 : (\mathbb{F}_2 : r(\varphi), a(\varphi), \delta_F(\varphi) = 1, \delta_{\varphi S}, v(\varphi)) &\xrightarrow{\mathbb{E}} (4; r(\psi) = r(\varphi) + 2, a(\psi) = a(\varphi) + 2, \delta_P = 0, \delta_\psi, \delta_{\psi P}); \\ D_4 : (\mathbb{F}_2 : r(\varphi), a(\varphi), \delta_F(\varphi) = 0, \delta_{\varphi S}, v(\varphi)) &\xrightarrow{\mathbb{E}} (4; r(\psi) = r(\varphi) + 2, a(\psi) = a(\varphi), \delta_P = 0, \delta_\psi, \delta_{\psi P}); \\ D_4 : (\mathbb{F}_2 : r(\tilde{\varphi}), a(\tilde{\varphi}), H(\tilde{\varphi}) = [S], \delta_{\tilde{\varphi} S}, v(\tilde{\varphi})) &\xrightarrow{\mathbb{E}} (4; r(\psi) = r(\tilde{\varphi}), a(\psi) = a(\tilde{\varphi}), \delta_P = 0, \delta_\psi, \delta_{\psi P}); \\ D_4 : (\mathbb{F}_2 : r(\tilde{\varphi}), a(\tilde{\varphi}), H(\tilde{\varphi}) = [F], \delta_{\tilde{\varphi} S}, v(\tilde{\varphi})) &\xrightarrow{\mathbb{E}} (4; r(\psi) = r(\tilde{\varphi}), a(\psi) = a(\tilde{\varphi}), \delta_P = 0, \delta_\psi, \delta_{\psi P}) \end{aligned}$$

where

$$\begin{aligned} \delta_\psi &= \begin{cases} 0 & \text{if } (\delta_{\varphi S} = \delta_{\tilde{\varphi} S}, r(\psi) \bmod 4) = (0, 2 \bmod 4) \\ 1 & \text{otherwise} \end{cases}, \\ \delta_{\psi P} &= \begin{cases} 0 & \text{if } (\delta_{\varphi S} = \delta_{\tilde{\varphi} S}, r(\psi) \bmod 4) = (0, 0 \bmod 4) \\ 1 & \text{otherwise} \end{cases}. \end{aligned}$$

Thus, similarly to Theorem 20 we obtain

Theorem 21. *All non-singular real quartics in $\mathbb{P}^3$ which are deformations of double coverings of real quadratic cone in $\mathbb{P}^3$ ramified in a non-singular degree 8 curve, and all isomorphism classes of these deformations are completely determined by the symbolical deformations (3.16), (3.17) and Figures 5, 6 and (3.7). For a fixed non-singular real quartic in $\mathbb{P}^3$ there are seven isomorphism classes of these deformations at most.*

Similar results for non-degenerate real quadrics in $\mathbb{P}^3$ (e. g. of hyperboloid type, or ellipsoid) had been obtained in [19]. Thus, Theorem 21 completes these results for a non-degenerate quadratic cone in $\mathbb{P}^3$.

All our results here (and results of [19] and [18] as well) are based on Global Torelli Theorem [20] and surjectivity of Torelli map [14] for K3 surfaces and some geometric and arithmetic considerations. Thus, our results are mainly existence and uniqueness results. It would be very interesting to construct exactly all representatives of connected components of moduli which we considered and classified. Our results about deformations reduce everything to construction of representatives of connected components of moduli of real non-singular

curves $A \in |-2K_{\mathbb{F}_m}|$, $0 \leq m \leq 4$, or degree 6 curves in $\mathbb{P}^2$. In many cases they are known by direct construction. E. g. see [7], [6], [8], [21], [24], [25], [26], [27], [28]. It would be very interesting to have such constructions for all connected components of moduli. It will make our results effective and useful for some questions which don't depend on deformation. This construction is surely possible because all our connected components of moduli are exactly labelled by some natural invariants, and number of the connected components is bounded.

REFERENCES

- [1] V.A. Alexeev, V.V. Nikulin, *Classification of del Pezzo surfaces with log-terminal singularities of index ≤ 2 , involutions of K3 surfaces, and reflection groups in Lobachevski spaces*, Doklady po matematike i prilozheniam. Matem. Institut im. Steklova. Akademia Nauk SSSR., **Vol. 2** (1988), no 2, 51–150 (in Russian). See English version in math.AG/0406536.
- [2] A.L. Besse, *Einstein Manifolds*, Ergeb. Math. Grenzgeb. (3), **10**, Springer-Verlag, 1987.
- [3] W. Barth, C. Peters, A. Van de Ven, *Compact Complex Surfaces*, Ergeb. Math. Grenzgeb. (3), **4**, Springer-Verlag, 1984.
- [4] A. Degtyarev, I. Itenberg, V. Kharlamov, *Real Enriques Surfaces*, Lect. Notes in Math., **1746** (2000), 259 pp.
- [5] S.K. Donaldson, *Yang–Mills invariants of smooth four-manifolds*, Geometry of low-dimensional manifolds (S.K. Donaldson, C.B. Thomas, eds.), Vol. 1. Cambridge University Press, Cambridge (1990), 5–40.
- [6] D.A. Gudkov, *On the topology of algebraic curves on a hyperboloid*, Usp. Mat. Nauk, **34** (1979), 26–32. English transl. in Russian Math. Surveys, **34** (1979), 27–35.
- [7] D.A. Gudkov, G.A. Utkin, *The topology of sixth-order curves and fourth order surfaces*, Gor'kov. Gos. Univ. Uchen. Zap., **87** (1969). English transl. in *Nine papers on Hilbert's 16 problem*, Amer. Math. Soc. Transl., **112** (1978), no 2.
- [8] D.A. Gudkov, E.I. Shustin, *A classification of nonsingular curves of order eight on an ellipsoid*, Methods of the Qualitative Theory of Differential Equations (E.A. Leontovich-Andronova, editor), Gor'kov. Gos. Univ., Gorki, (1980), 104–107 (in Russian).
- [9] Daniel Huybrechts, *Moduli spaces of hyperkähler manifolds and mirror symmetry*, math.AG/0210219 v3.
- [10] I. V. Itenberg, *Curves of degree 6 with one non-degenerate double point and groups of monodromy of non-singular curves*, Real Algebraic Geometry, Proceedings, Rennes 1991, Lecture Notes in Math., **1524**, Springer, (1992), 267–288.
- [11] I. V. Itenberg, *Rigid isotopy classification of curves of degree 6 with one nondegenerate double point*, Topology of Manifolds and Varieties, Advances in Soviet Math., **18** (1994), 193–208.
- [12] V.M. Kharlamov, *Additional congruences for the Euler characteristic of real algebraic manifolds of even dimensions*, Funkt. Anal. Prilozhen., **9** (1975), 51–60. English transl. in Funct. Anal. Appl., **9** (1975), 134–141.
- [13] V.M. Kharlamov, *The topological types of nonsingular surfaces in $\mathbb{RP}^3$ of degree four*, Funkt. Anal. Prilozhen., **10** (1976), 55–68. English transl. in Funct. Anal. Appl., **10** (1976), 295–305.
- [14] V.S. Kulikov, *Degenerations of K3 surfaces and Enriques surfaces*, Izv. AN SSSR. Ser. Mat., **41** (1977), 1008–1042. English transl. in Math. USSR Izv., **11** (1977), 957–989.
- [15] V.V. Nikulin, *Integral symmetric bilinear forms and some of their geometric applications*, Izv. Akad. Nauk SSSR. Ser. Mat., **43** (1979), 111–177. English transl. in Math. USSR Izv., **14** (1980), 103–167.
- [16] V.V. Nikulin, *On the quotient groups of the automorphism groups of hyperbolic forms by the subgroups generated by 2-reflections*, Algebraic-geometric applications, Current Problems in Math., Vsesoyuz. Inst. Nauch. i Tekhn. Informatsii, Moscow, **18** (1981), 3–114. English transl. in J. Soviet Math., **22** (1983), 1401–1476.

- [17] V.V. Nikulin, *Involutions of integral quadratic forms and their applications to real algebraic geometry*, Izv. Akad. Nauk SSSR Ser. Mat., **47** (1983), 109–188. English transl. in Math. USSR Izv., **22** (1984), 99–172.
- [18] V.V. Nikulin, *Remarks on connected components of moduli of real polarized K3 surfaces*, Preprint math.AG/0507197, 22 pages.
- [19] V.V. Nikulin, S. Saito, *Real K3 surfaces with non-symplectic involution and applications*, Proc. London Math. Soc., **90** (2005), 591–654. (See also math.AG/0312396)
- [20] I.I. Piateckii-Shapiro, I.R. Shafarevich, *A Torelli theorem for algebraic surfaces of type K3*, **35** (1971), 530–572. English transl. in Math. USSR Izvestija, **5** (1971), 547–588.
- [21] V.A. Rokhlin, *Complex topological characteristics of real algebraic curves*, Usp. Mat. Nauk, **33** (1978), 77–89. English transl. in Russian Math. Surveys, **33** (1978), 85–98.
- [22] B. Saint-Donat, *Projective models of K-3 surfaces*, Amer. J. Math., **96** (1974), 602–639.
- [23] I.R. Shafarevich et al, *Algebraic surfaces*, Trudy Matem. Inst. Steklov, **75** (1965). English transl. in Proc. Steklov Inst. Math., **75** (1965).
- [24] O.Ya. Viro, *Curves of degree 7, curves of degree 8 and the Ragsdale conjecture*, Dokl. Akad. Nauk SSSR, **254** (1980), 1306–1310. English transl. in Soviet. Math. Dokl., **22** (1980), 566–570.
- [25] G. Wilson, *Hilbert's sixteenth problem*, Topology, **17** (1978), 53–73.
- [26] V.I. Zvonilov, *Complex topological characteristics of real algebraic curves on surfaces*, Funkt. Anal. i Pril., **16** (1982), 56–57. English transl. in Funct. Anal. Appl., **16** (1982), 202–203.
- [27] V.I. Zvonilov, *Complex orientations of real algebraic curves with singularities*, Dokl. AN SSSR, **268** (1983), 22–26. English transl. in Soviet Math. Dokl., **27** (1983), 14–17.
- [28] V.I. Zvonilov, *Complex topological invariants of real algebraic curves on a hyperboloid and on an ellipsoid*, Algebra i Analiz, **3** (1992), 88–108. English transl. in St. Petersburg Math. J., **3** (1992), 1023–1042.

DEPARTMENT OF PURE MATHEMATICS, THE UNIVERSITY OF LIVERPOOL, LIVERPOOL, L69 3BX,
UNITED KINGDOM, STEKLOV MATHEMATICAL INSTITUTE, UL. GUBKINA 8, MOSCOW, GSP-1, RUSSIA
E-mail address: vnikulin@liv.ac.uk, vvnikulin@list.ru

DEPARTMENT OF MATHEMATICS EDUCATION, HAKODATE CAMPUS, HOKKAIDO UNIVERSITY OF EDUCATION,
1-2 HACHIMAN-CHO, HAKODATE 040-8567, JAPAN
E-mail address: sachi63@cc.hokkyodai.ac.jp